

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :					
--	--	--	--	--	--

(34) International Patent Classification:	A61K 38/00, C07K 100, C07H 21/02, 21/04	A1
(11) International Publication Number:	WO 98/55132	
(43) International Publication Date:	10 December 1998 (10.12.98)	

(21) International Application Number: PCT/US98/10104

(22) International Filing Date: 18 May 1998 (18.05.98)

(31) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GR, GH, GM, GW, HT, IN, IT, JP, KE, KG, KH, KR, KZ, LA, LG, LI, LU, LV, LY, MA, MD, MG, MK, MN, MU, MV, MW, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SD, SG, SI, SK, SL, SM, SN, SR, ST, SV, SZ, TD, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VE, VN, YU, ZA, ZM, ZW

(32) Priority Date(s): 1997-05-15 (US 08/658,100)

(33) Priority Application Number(s): 08/658,100

(34) Priority Filing Date(s): 1997-05-15

(35) Priority Claimed From:

(36) Priority Claimed To:

(40) Publication Date: 1999-01-28

(41) Publication Kind: Full Text

(42) Publication Language: English

(43) Publication Date of Abstract: 1999-01-28

(44) Publication Date of Claims: 1999-01-28

(45) Publication Date of Description: 1999-01-28

(46) Publication Date of Figures: 1999-01-28

(47) Publication Date of Tables: 1999-01-28

(48) Publication Date of Drawings: 1999-01-28

(49) Publication Date of Other Material: 1999-01-28

(50) Publication Date of Other Material: 1999-01-28

(51) International Classification: G06F 01/00

(52) International Classification: G06F 01/00

(53) International Classification: G06F 01/00

(54) International Classification: G06F 01/00

(55) International Classification: G06F 01/00

(56) International Classification: G06F 01/00

(57) International Classification: G06F 01/00

(58) International Classification: G06F 01/00

(59) International Classification: G06F 01/00

(60) International Classification: G06F 01/00

(61) International Classification: G06F 01/00

(62) International Classification: G06F 01/00

(63) International Classification: G06F 01/00

(64) International Classification: G06F 01/00

(65) International Classification: G06F 01/00

(66) International Classification: G06F 01/00

(67) International Classification: G06F 01/00

(68) International Classification: G06F 01/00

(69) International Classification: G06F 01/00

(70) International Classification: G06F 01/00

(71) International Classification: G06F 01/00

(72) International Classification: G06F 01/00

(73) International Classification: G06F 01/00

(74) International Classification: G06F 01/00

(75) International Classification: G06F 01/00

(76) International Classification: G06F 01/00

(77) International Classification: G06F 01/00

(78) International Classification: G06F 01/00

(79) International Classification: G06F 01/00

(80) International Classification: G06F 01/00

(81) International Classification: G06F 01/00

(82) International Classification: G06F 01/00

(83) International Classification: G06F 01/00

(84) International Classification: G06F 01/00

(85) International Classification: G06F 01/00

(86) International Classification: G06F 01/00

(87) International Classification: G06F 01/00

(88) International Classification: G06F 01/00

(89) International Classification: G06F 01/00

(90) International Classification: G06F 01/00

(91) International Classification: G06F 01/00

(92) International Classification: G06F 01/00

(93) International Classification: G06F 01/00

(94) International Classification: G06F 01/00

(95) International Classification: G06F 01/00

(96) International Classification: G06F 01/00

(97) International Classification: G06F 01/00

(98) International Classification: G06F 01/00

(99) International Classification: G06F 01/00

(100) International Classification: G06F 01/00

(30) Priority Data:	
08/683,162	US
09/076,606	US
2 June 1997 (02.06.97)	US
12 May 1998 (12.05.98)	US

71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA (USUS); 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US).

72) Investors: PRUSNER, Stanley, B.; 400 Picacho Street, San Francisco, CA 94116 (US); COHEN, Fred, R.; 767 Rhode Island Street, San Francisco, CA 94107 (US); JAMES, Thomas, L.; 33 Camino Margatta, Nicasio, CA 94946 (US).

154) Agent: BOZICHVIC, Kart; Bozichvic & Reed LLP, Suite 200,
283 Hamilton Avenue, Palo Alto, CA 94301 (US).

44) Title: PRION PROTEIN MODULATOR FACTOR (PPMF) AND PRION RESISTANT ANIMALS

Abstract

[illegible]

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LT	Lithuania	EE	Estonia
AM	Armenia	FI	Finland	LU	Luxembourg	IE	Ireland
AN	Andorra	FR	France	LJ	Liechtenstein	IS	Iceland
AO	Angola	GB	Great Britain	LV	Latvia	IT	Italy
AT	Austria	GR	Greece	LI	Liechtenstein	JO	Jordan
BA	Bosnia and Herzegovina	GU	Guam	MD	Moldova	KZ	Kazakhstan
BB	Barbados	HK	Hong Kong	MG	Madagascar	KG	Kyrgyzstan
BD	Bangladesh	HN	Honduras	ML	Malawi	LA	Laos
BE	Belgium	IE	Ireland	MP	Marshall Islands	LC	Liechtenstein
BF	Burkina Faso	IL	Israel	MR	Moldova	LI	Liechtenstein
BG	Bulgaria	IT	Italy	MT	Malta	LV	Latvia
BJ	Benin	JP	Japan	NI	Nicaragua	LT	Lithuania
BK	Bosnia and Herzegovina	KE	Kenya	NO	Norway	LU	Luxembourg
BM	Bermuda	KG	Kazakhstan	NZ	New Zealand	UK	United Kingdom
BN	Brunei	LA	Laos	PL	Poland	US	United States of America
BO	Bolivia	LV	Latvia	PT	Portugal	VI	Virgin Islands
BR	Brazil	MY	Malaysia	RO	Romania	VG	Virgin Islands
BS	Bahamas	MX	Mexico	RU	Russia	VI	Virgin Islands
BT	Bhutan	NC	New Caledonia	SD	Sudan	WV	West Virginia
BV	Bouvet Island	NE	Netherlands	SE	Sweden	WY	Wyoming
BZ	Belize	NI	Nicaragua	SG	Singapore	DE	Germany
CA	Canada	NP	Nepal	SI	Slovenia	DK	Denmark
CC	Cocos (Keeling) Islands	NR	Nauru	SK	Slovakia	DO	Dominican Republic
CD	Congo	NU	Niue	SM	San Marino	EC	Ecuador
CF	Central African Republic	OM	Oman	SN	Senegal	FR	France
CG	Congo	PA	Panama	SR	Suriname	GR	Greece
CH	China	PE	Peru	SV	El Salvador	DE	Germany
CI	Cote d'Ivoire	PH	Philippines	TD	Chad	GI	Gibraltar
CK	Cook Islands	PK	Pakistan	TH	Thailand	GN	Guinea
CL	Chile	PL	Poland	TM	Turkmenistan	GW	Guinea-Bissau
CM	Camerun	PR	People's Republic of China	TR	Turkey	HT	Haiti
CN	China	PS	Palestine	TV	Tuvalu	HR	Croatia
CO	Colombia	QA	Qatar	UG	Uganda	HU	Hungary
CZ	Czech Republic	RE	Reunion	US	United States of America	UA	Ukraine
CU	Cuba	RO	Romania	UY	Uruguay	UA	Ukraine
CV	Cape Verde	RU	Russia	UZ	Uzbekistan	NP	Nepal
CY	Cyprus	RS	Russia	VE	Venezuela	NI	Nicaragua
DE	Germany	RU	Russia	VC	St. Vincent and the Grenadines	NR	Nauru
DM	Dominica	SE	Sweden	VG	Virgin Islands	NC	New Caledonia
DO	Dominican Republic	SI	Slovenia	VG	Virgin Islands	NE	Netherlands
EC	Ecuador	LI	Liechtenstein	VG	Virgin Islands	NG	Nigeria
EE	Estonia	LU	Luxembourg	VG	Virgin Islands	NI	Nicaragua
EG	Egypt	LV	Latvia	VG	Virgin Islands	NL	Netherlands
EH	Western Sahara	LT	Lithuania	VG	Virgin Islands	NO	Norway
ER	Eritrea	LU	Luxembourg	VG	Virgin Islands	NZ	New Zealand
ES	Spain	LV	Latvia	VG	Virgin Islands	PL	Poland
EU	European Union	LV	Latvia	VG	Virgin Islands	PT	Portugal
EV	Evros	LV	Latvia	VG	Virgin Islands	RO	Romania
EW	Western Sahara	LV	Latvia	VG	Virgin Islands	RU	Russia
EX	Excluded Territories	LV	Latvia	VG	Virgin Islands	SD	Sudan
FJ	Fiji	LV	Latvia	VG	Virgin Islands	SE	Sweden
FK	Falkland Islands	LV	Latvia	VG	Virgin Islands	SG	Singapore
FM	Federated States of Micronesia	LV	Latvia	VG	Virgin Islands	SI	Slovenia
FO	Faroese Islands	LV	Latvia	VG	Virgin Islands	SK	Slovakia
FR	France	LV	Latvia	VG	Virgin Islands	SM	San Marino
FS	French Southern Territories	LV	Latvia	VG	Virgin Islands	SN	Senegal
GA	Gabon	LV	Latvia	VG	Virgin Islands	SR	Suriname
GB	Great Britain	LV	Latvia	VG	Virgin Islands	SV	El Salvador
GD	Grenada	LV	Latvia	VG	Virgin Islands	TD	Chad
GE	Georgia	LV	Latvia	VG	Virgin Islands	TH	Thailand
GF	French Guiana	LV	Latvia	VG	Virgin Islands	TM	Turkmenistan
GG	Guernsey	LV	Latvia	VG	Virgin Islands	TR	Turkey
GH	Ghana	LV	Latvia	VG	Virgin Islands	TV	Tuvalu
GI	Gibraltar	LV	Latvia	VG	Virgin Islands	UG	Uganda
GL	Greenland	LV	Latvia	VG	Virgin Islands	UY	Uruguay
GM	Gambia	LV	Latvia	VG	Virgin Islands	UZ	Uzbekistan
GN	Guinea	LV	Latvia	VG	Virgin Islands	VE	Venezuela
GO	Goa	LV	Latvia	VG	Virgin Islands	VC	St. Vincent and the Grenadines
GP	Guadeloupe	LV	Latvia	VG	Virgin Islands	VG	Virgin Islands

PRION PROTEIN MODULATOR FACTOR (PPMF) AND PRION RESISTANT ANIMALS

FIELD OF THE INVENTION

This invention relates generally to proteins, functionally equivalent pharmacophores and methods of creating a gene to make prion resistant animals. More specifically, this invention relates to (1) a discontinuous epitope of PrP protein, (2) prion protein modulator factor (PPMF) which binds the epitope, (3) functional pharmacophores of (1) and (2) and (4) prion resistant animals and their PrP genes.

BACKGROUND OF THE INVENTION

Prions are infectious pathogens that cause central nervous system spongiform encephalopathies in humans and animals. Prions are distinct from bacteria, viruses and viroids. The predominant hypothesis at present is that no nucleic acid component is necessary for infectivity of prion protein. Further, a prion which infects one species of animal (e.g., a human) will not infect another (e.g., a mouse).

A major step in the study of prions and the diseases that they cause was the discovery and purification of a protein designated prion protein ("PrP") [Bolton et al., *Science* 218:1309-11 (1982); Prusiner et al., *Biochemistry* 21:6942-50 (1982); McKinley et al., *Cell* 35:57-62 (1983)]. Complete prion protein-encoding genes have since been cloned, sequenced and expressed in transgenic animals. PrP^{Sc} is encoded by a single-copy host gene [Basler et al., *Cell* 46:417-28 (1986)] and is normally found at the outer surface of neurons. A leading hypothesis is that prion diseases result from conversion of PrP^C into a modified form called PrP^{Sc}.

It appears that the scrapie isoform of the prion protein (PrP^{Sc}) is necessary for both the transmission and pathogenesis of the transmissible neurodegenerative diseases of animals and humans. See Prusiner, S.B., "Molecular biology of prion disease," *Science* 252:1515-1522 (1991). The most common prion diseases of animals are scrapie of sheep and goats and bovine spongiform encephalopathy (BSE) of cattle [Wilesmith, J. and Wells, *Microbiol. Immunol.* 17:221-38 (1991)]. Four prion diseases of humans have been identified: (1) kuru,

(2) Creutzfeldt-Jacob Disease (CJD), (3) Gerstmann-Sträussler-Scheinker Disease (GSS), and (4) fatal familial insomnia (FFI) [Gajdusek, D.C., *Science* 197:943-960 (1977); Medori et al., *N. Engl. J. Med.* 326:444-449 (1992)]. The presentation of human prion diseases as sporadic, genetic and infectious illnesses initially posed a conundrum which has been explained by the cellular genetic origin of PrP.

The initial transmission of CJD to experimental primates has a rich history beginning with William Hadlow's recognition of the similarity between kuru and scrapie. In 1959, Hadlow suggested that extracts prepared from patients dying of kuru be inoculated into nonhuman primates and that the animals be observed for disease that was predicted to occur after a prolonged incubation period [Hadlow, W.J., *Lancet* 2:289-290 (1959)]. Seven years later, Gajdusek, Gibbs and Alpers demonstrated the transmissibility of kuru to chimpanzees after incubation periods ranging from 18 to 21 months [Gajdusek et al., *Nature* 209:794-796 (1966)]. The similarity of the neuropathology of kuru with that of CJD [Kluwe et al., *Lab Invest.* 8:799-847 (1959)] prompted similar experiments with chimpanzees and transmissions of disease were reported in 1968 [Gibbs, Jr. et al., *Science* 161:388-389 (1968)]. Over the last 25 years, about 300 cases of CJD, kuru and GSS have been transmitted to a variety of apes and monkeys.

The expense, scarcity and often perceived inhumanity of such experiments have restricted this work and thus limited the accumulation of knowledge. While the most reliable transmission data has been said to emanate from studies using nonhuman primates, some cases of human prion disease have been transmitted to rodents but apparently with less regularity [Gibbs, Jr. et al., *Slow Transmissible Diseases of the Nervous System*, Vol. 2, S.B. Prusiner and W.J. Hadlow, eds (New York: Academic Press), pp. 87-110 (1979); Tateishi et al., *Prion Diseases of Humans and Animals*, Prusiner et al., eds (London: Ellis Horwood), pp. 129-134 (1992)].

The infrequent transmission of human prion disease to rodents has been cited as an example of the "species barrier" first described by Pattison in his studies of passing the scrapie agent between sheep and rodents [Pattison, I.H., *NINDB Monograph* 2, D.C. Gajdusek, C.J. Gibbs Jr. and M.P. Alpers, eds (Washington, D.C.: U.S. Government Printing), pp. 249-257 (1965)]. In those investigations, the initial passage of prions from one species to another was associated with a prolonged incubation time with only a few animals

developing illness. Subsequent passage in the same species was characterized by all the animals becoming ill after greatly shortened incubation times.

The molecular basis for the species barrier between Syrian hamster (SHA) and mouse was shown to reside in the sequence of the PrP gene using transgenic (Tg) mice [Scott et al., *Cell* 59:847-857 (1989)]. SHA PrP differs from Mo PrP at 16 positions out of 254 amino acid residues [Bastier et al., *Cell* 46:417-428 (1986); Loch et al., *Proc. Natl. Acad. Sci. USA* 83:6372-6376 (1986)]. Tg(SHA PrP) mice expressing SHA PrP had abbreviated incubation times when inoculated with SHA prions. When similar studies were performed

with mice expressing the human or ovine PrP transgenes, the species barrier was not abrogated, i.e., the percentage of animals which became infected were unacceptably low and the incubation times were unacceptably long. Thus, it has not been possible, for example in the case of human prions, to use transgenic animals (such as mice containing a PrP gene of another species) to reliably test a sample to determine if that sample is infected with prions. The seriousness of the health risk resulting from the lack of such a test is exemplified below.

More than 45 young adults previously treated with HGH derived from human pituitaries have developed CJD [Koch et al., *N. Engl. J. Med.* 313:731-733 (1985); Brown et al., *Lancet* 340:24-27 (1992); Fradkin et al., *JAMA* 265:880-884 (1991); Buchanan et al., *Br. Med. J.* 302:824-828 (1991)]. Fortunately, recombinant HGH is now used, although the seemingly remote possibility has been raised that increased expression of wt PrP^C stimulated by high HGH might induce prion disease [Lamezas et al., *Biochem. Biophys. Res. Commun.* 196:1163-1169 (1993)]. That the HGH prepared from pituitaries was contaminated with prions is supported by the transmission of prion disease to a monkey 66

months after inoculation with a suspect lot of HGH [Gibbs, Jr. et al., *N. Engl. J. Med.* 328:358-359 (1993)]. The long incubation times associated with prion diseases will not reveal the full extent of iatrogenic CJD for decades in thousands of people treated with HGH worldwide. Iatrogenic CJD also appears to have developed in four infertile women treated with contaminated human pituitary-derived gonadotrophin hormone [Healy et al., *Br. J. Med.* 307:517-518 (1993); Cochius et al., *Aust. N.Z. J. Med.* 20:592-593 (1990); Cochius et al., *J. Neurol. Neurosurg. Psychiatry* 55:1094-1095 (1992)] as well as at least 11 patients receiving dura mater grafts [Nisbet et al., *J. Am. Med. Assoc.* 261:1118 (1989); Thadani et al., *J. Neurosurg.* 69:766-769 (1988); Willison et al., *J. Neurosurg. Psychiatry* 54:940

(1991); Brown et al., *Lancet* 340:24-27 (1992)]. These cases of iatrogenic CJD underscore the need for screening pharmaceuticals that might possibly be contaminated with prions.

Such a method is provided in U.S. Patent 5,565,186, issued October 15, 1996, which provides for inoculating transgenic mice with a sample suspected of containing prions and waiting to see if the mice develop symptoms of prion disease. The present invention provides a protein designated PPMF which has a range of uses including facilitating the speed and sensitivity of assays which use transgenic animals to detect prions.

SUMMARY OF THE INVENTION

Prion Protein Modulator Factor (PPMF) is disclosed which protein is characterized by binding to PrP^C and facilitating a conformation change from PrP^C to PrP^{Sc}. The protein PPMF is species specific meaning that PPMF from one species of mammal will only bind to PrP^C of the same or a genetically similar species. PPMF binds to a discontinuous epitope of PrP protein and the epitope is also disclosed in terms of its amino acid sequence but, more importantly, in terms of its 3-dimensional configuration which defines the negative image of the binding surface of PPMF. Using the 3-dimensional coordinates defining the binding surface of both PPMF and the discontinuous epitope of PrP protein, functionally equivalent pharmacophores are disclosed and described as are methods of generating such pharmacophores. PPMF, its binding epitope on PrP protein and functionally equivalent pharmacophores of each are useful in assays and as therapeutic agents. By identifying the PrP^C binding epitope of PPMF, PrP genes can be created which render an animal resistant to prion disease.

An aspect of the invention is the 3-dimensionally defined discontinuous epitope of PrP protein and functionally equivalent pharmacophores.

Another aspect of the invention is PPMF and functionally equivalent pharmacophores with a 3-dimensionally defined surface in the negative image of the discontinuous epitope of PrP protein.

Another aspect of the invention is a method of identifying compounds which inhibit PrP^{Sc} formation which method comprises (1) identifying a discontinuous PPMF binding site on PrP^C; (2) matching compounds in a library with the binding site regarding spatial orientation; (3) screening candidate compounds visually using computer generated molecular

display software; and (4) screening actual compounds against PrP^C in the presence of PrP^{Sc} to determine compounds which inhibit conversion of PrP^C to PrP^{Sc}.

Another aspect of the invention is a PrP gene with codon substitutions made at position which encode amino acids of the discontinuous epitope, in which substituents are made to encode a protein which (1) binds PPMF more readily than native PrP or (2) does not bind PPMF, thereby providing genes which can provide transgenic animals which are resistant to prion infection.

Another aspect of the invention comprises administering PPMF to a transgenic animal with a chimeric artificial PrP gene comprising portions of the host animal, e.g. and portions of a host animal and a middle portion of a genetically diverse test animal, wherein the middle portion includes specific alterations designed to match that of a disease state of such a host. The PPMF can be administered to a transgenic animal containing a chimeric gene in order to reduce the time for prion disease to be exhibited.

The artificial gene includes a sequence such that when inserted into the genome of an animal (such as a mouse), it renders the animal susceptible to infection with prions which normally would infect only a specific species of genetically diverse animal (such as a human, cow, sheep, pig, chicken, cat or dog). The artificial PrP gene may be comprised partially or completely of an artificial polynucleotide sequence, i.e. codon sequences not present in any native PrP gene sequence. Alternatively, the artificial gene may be comprised of the codon sequence of a host animal with one or more codon substitutions being made wherein the substitutions are preferably corresponding PrP gene codons from a genetically diverse animal, meaning that PrP gene differs from the PrP gene of the host animal by 20 or more codons. Transgenic animals containing elevated levels of expression of the PrP gene which can be obtained, for example, by over expression of the gene with an enhanced promoter and/or with high copy numbers of the natural PrP gene of a genetically diverse animal are also disclosed. Hybrid transgenic animals include animals resulting from a cross between two transgenic animals and in particular a cross between a transgenic animal containing the entire prion protein gene of a genetically diverse animal (e.g., a mouse containing a human prion protein gene) and an animal with its endogenous prion protein gene disrupted (e.g., a mouse with an ablated prion protein gene). Hybrids also specifically include crossing a transgenic animal having a chimeric prion protein gene with an animal with its endogenous prion protein gene ablated.

PPMF or ligands which provide a surface that fits 3-dimensionally defined pharmacophores of the invention can be administered to transgenic animals which, due to their genetic make up will develop disease from inoculation with prions which would generally only infect a genetically diverse animal, e.g., a transgenic mouse will consistently become infected with prions which generally will only infect a human and symptoms of the infection will become apparent in a short period e.g., 100 days or less. The PPMF and/or pharmacophores are administered to transgenic animals that are used in assays to test samples of any given material to determine if the material includes prions which would infect another animal (such as a human) if the material were ingested or injected.

Standardized prion preparations may be used to inoculate animals to create controls when carrying out an assay of the invention. The standardized prion preparation will always contain prions which will infect a genetically modified animal that will develop clinical signs of CNS dysfunction within a set period of time.

An aspect of the invention is a discontinuous epitope of PrP protein which is precisely defined in 3-dimensional space by distances and angles between amino acids making up the surface of the 3-dimensional epitope and functionally equivalent pharmacophores defining the same 3-dimensional space.

Another aspect of the invention is the binding site of PPMF to PrP which is defined by the negative image of the defined 3-dimensional space of the epitope.

An object of the invention is to provide an isolated protein designated Prion Protein Modulator Factor (PPMF) which is characterized by binding to PrP^C and facilitating a conformational change to PrP^{Sc}.

Another object of the invention is to provide transgenic animals such as cows and sheep which have their PrP^C gene modified to express PrP^C protein with a modified epitope for binding PPMF in a manner which renders the animal resistant to prion diseases.

Another object is to provide antibodies and pharmacophores which bind to PPMF (or epitopes of PrP^C which bind PPMF) which molecules block PPMF - PrP^C binding and thereby act as a therapeutic for prion disease.

Another object of the invention is to provide for a method of testing samples for the presence of prions which preferably involves creating two groups of nonhuman mammals with their genome altered so that they are susceptible to infection with prions which generally only infect a genetically diverse animal. The first group of animals is infected with

5 WO 98/51332 PCT/US98/01004

A test sample while being given PPMF and the second group is infected with a standardized prion preparation while being given PPMF. Both groups of mammals are observed and the presence of prions in the sample can be deduced if the first group of animals develop symptoms of prion infection. The time needed to develop symptoms is reduced by the PPMF.

10 An advantage of the invention is that the PPMF and pharmacophores thereof reduce the time needed for and increase the sensitivity of an assay.

A feature of the invention is that the PPMF and pharmacophores thereof do not bind to PrP^{Sc}.

15 Another advantage is that the transgenic and hybrid animals injected with PPMF can quickly (100 days or less) detect prions in a sample at very low levels, e.g., 1 part per million, and even as low as 1 part per billion.

Still another advantage is that the transgenic and hybrid animals injected with PPMF can be used for an assay which is highly accurate, i.e., does not provide false positives and consistently determines the presence of prions.

Another object of the invention is to provide an assay wherein a sample is assayed for the presence of a PrP^{Sc} by adding to the sample PrP^C in combination with PPMF, a pharmacophore thereof or an antibody of PrP^C which binds to PrP^C at its PrP^C/PPMF binding epitope whereby the assay amplifies the amount of PrP^{Sc} (if any) present in the sample.

20 These and other objects, advantages, and features of the invention will become apparent to those persons skilled in the art upon reading the details of the chimeric gene, assay method, and transgenic mouse as more fully described below.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figure 1 is a schematic representation of how PPMF converts PrP^C to PrP^{Sc}.

Figure 2 is a schematic representation of PrP^C showing the binding epitope for PPMF.

Figure 3 shows a comparison of a number of PrP sequences with a segment of mouse PrP.

30 Figure 4 shows the 3-dimensional geometries of the discontinuous binding site of PrP^{Sc}.

Figure 5 is a computer generated comparison of the ten best-scoring structures of rPrP shown with a best-fit superposition of backbone atoms for residues 113-227 (stereoview). In Figures 10, 12 and 14, the color scheme is: disulfide between Cys¹⁷⁹ and Cys¹⁸¹, yellow; sites of glycosylation in PrP^C, i.e., Asn¹⁴¹ and Asn¹⁹⁷, gold; hydrophobic

5 globule composed of residues 113-126, red; helices, pink; loops, gray; residues 129-134, green, encompassing strand S2; the arrows span residues 129-131 and 161-163, as these show a closer resemblance to β -sheet. The structures were generated with the program DIANA [Gunter, et al. J. Mol. Biol. 217, 517-530 (1990)] followed by energy minimization with AMBER 4.1 [Pearlman, et al. AMBER 4.1 (UCSF 1995)]. Structure generation

10 parameters are as follows: 2401 distant restraints (intrareidue, 858; sequential (i-i+1), 753; i-i+2, 195; i-i+3, 233; i-i+4, 109; and i-i+5, 253 for amino acid; hydrogen bond restraints, 44; distance restraints violations $\times 0.5A$ per structure, 30; AMBER energy, -1443 \pm 111 kcal/mol. Precision of structures: atomic RMSD for all backbone heavy atoms of residues 128-227, $\pm 1.9A$. The distance restraint violations and precision in some molecular

15 moieties reflect the conformational heterogeneity of rPrP.

Figures 6A and 6B show a portion of the 3D ¹³C-NOESY spectrum corresponding to ¹³C planes of the unresolved Val¹⁴⁶ methyl resonances and the Ser²²⁷ resonances. The diagonal peaks and mirrored cross-peaks for each ¹³C-¹³C connectivity is shown. The solid lines connecting peaks designate NOE connectivities.

Figure 7 is a computer generated structure of the interaction of the hydrophobic globule, with van der Waals rendering the atoms in residues 113-127, with the first β -strand.

Figure 8 is a computer generated structure showing van der Waals surface of PrP turned 90 from Figure 5, illustrating the interaction of helix A with helix C. Helices A, B and C are respectively colored magenta, white and gold.

20 Figure 9 is a computer generated stereoview highlighting in white the residues corresponding to point mutations leading to human prion diseases.

Figures 5, 7, 8 and 9 were generated with Midaspas.

Coordinates are deposited with the Brookhaven Protein Data Bank.

Figure 10 shows the secondary structure diagram of rPrP. NOE connectivities are denoted by lines, where the thickness qualitatively represents the relative intensity (weak, medium or strong) of the NOE cross-peaks, and i designates the residue number for rPrP.

30 $d_{NH}(i, i+3)$ denotes an NOE between the α -proton of residue i and the amide proton of $d_{NH}(i, i+3)$.

residue i+3. The long-range NOE line indicates by height the relative number of NOE cross-peaks between residues i-i+4. The α -carbon chemical shift index with contiguous up bars designating α -helix and down bars designating β -strand. Regions of secondary structure are depicted by helices for α -helices and broad arrows for β -strands. Hydrogen exchange is calculated from the intensity of proton NOE cross-peak between the amide and water: open circles for slow, filled for fast, and half-filled for medium exchange rate. No circle indicates spectral overlap or proline. The secondary structure diagram was created using the program Vinct[®] (E. La Rosa, A. Stern, J. Hoch, Copyright 1996, Rowland Institute for Science).

Figure 11 schematically shows critical epitope positions for hamster, mouse, human and sheep PrP proteins and relates such to their position on the PrP molecule.

Figures 12-15 are flow diagrams of a specific embodiment of the assay method of the invention. Figure 12 is a flow diagram [showing the invention applied to the PPMF binding site] of syrian hamster PrP^{Sc} molecule. Figure 13 is a flow diagram [showing the invention applied to the PPMF binding site] for the mouse PPMF binding site. Figure 14 is a flow diagram [showing the invention applied to the PPMF binding site] for the sheep PPMF binding site. Figure 15 is a flow diagram [showing the invention applied to the PPMF binding site] for the human PPMF binding site.

Figure 16 is a schematic diagram of compounds which inhibit PrP^{Sc} formation showing where and how compounds bind a PPMF binding epitope on PrP^{Sc}.

Figure 17 is a computer generated schematic diagram of rPrP(90-231) showing the residues involved in the discontinuous epitope to which PPMF binds.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Before the present proteins, pharmacophores, assay methodology, and transgenic and hybrid animals used in the assay are described, it is to be understood that this invention is not limited to particular protein, pharmacophores, assay methods, chimeric and artificial genes, prion preparation or transgenic and hybrid animals described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention

belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

DEFINITIONS

The term "PPMF" is used for Prion Protein Modulator Factor which is a protein which can be glycosylated and is characterized by binding to PrP^{Sc} and facilitating a conformational change from PrP^C to PrP^{Sc}. The term encompasses any PPMF from any animal allowing for specific differences between different species of animals. The PPMF compounds of the present invention are more particularly characterized herein.

A pharmacophore is a compound that has a specific biochemical activity which activity is obtained by the 3-dimensional physical shape of the compound and the electrochemical properties of the atoms making up the compound. Specific pharmacophores of the invention mimic (1) the discontinuous epitope on PrP^C to which PPMF binds or (2) the surface of PPMF which bind to PrP^{Sc}. Thus a pharmacophores of the invention have a shape (i.e., the geometric specifications) substantially as defined by Figure 4 or the negative image thereof. The term mimetic is used here interchangeably with pharmacophore and covers peptides, and small molecules which mimic the shape and spatial positioning of the functional groups on the molecule being mimicked.

The terms "treatment", "treating" and "treat" and the like are used herein to generally mean obtaining a desired pharmacological and/or physiological effect. The effect may be prophylactic in terms of completely or partially preventing a prion disease or symptom thereof and/or may be therapeutic in terms of partially or completely curing a prion disease or adverse effect attributable to the disease. The "treatment" as used herein covers any

WO 9855132
PCT/US98/10104
accidental infection of children with human prions from contaminated preparations of human growth hormone.

The term "Familial CJD" refers to a form of CJD which occurs rarely in families and is inevitably caused by mutations of the human prion protein gene. The disease results from an autosomal dominant disorder. Family members who inherit the mutations succumb to CJD.

The term "Gerstmann-Strassler-Scheinker Disease" abbreviated as "GSS" refers to a form of inherited human prion disease. The disease occurs from an autosomal dominant disorder. Family members who inherit the mutant gene succumb to GSS.

The term "prion" shall mean an infectious particle known to cause diseases (spongiform encephalopathies) in humans and animals. The term "prion" is a contraction of the words "protein" and "infection" and the particles are comprised largely if not exclusively of PrP^{Sc} molecules encoded by a PrP gene which expresses PrP^C which changes conformation to become PrP^{Sc}. Prions are distinct from bacteria, viruses and viroids.

Known prions include those which infect animals to cause scrapie, a transmissible, degenerative disease of the nervous system of sheep and goats as well as bovine spongiform encephalopathies (BSE) or mad cow disease and feline spongiform encephalopathies of cats. Four prion diseases known to affect humans are (1) kuru, (2) Creutzfeldt-Jakob Disease (CJD), (3) Gerstmann-Strassler-Scheinker Disease (GSS), and (4) fatal familial insomnia (FFI). As used herein prion includes all forms of prions causing all or any of these diseases or others in any animals used - and in particular in humans and in domesticated farm animals.

The terms "PrP gene" and "prion protein gene" are used interchangeably herein to describe genetic material which expresses PrP proteins, including proteins with polymorphisms and mutations such as those listed herein under the subheading "Pathogenic Mutations and Polymorphisms." Unless stated otherwise the term refers to the native wild-type gene and not to an artificially altered gene. The PrP gene can be from any animal including the "host" and "test" animals described herein and any and all polymorphisms and mutations thereof; it being recognized that the terms include other such PrP genes that are yet to be discovered. The term "PrP gene" refers generally to any gene of any species which encodes any form of a PrP amino acid sequences including any prion protein. Some commonly known PrP sequences are described in Gabriel et al., *Proc. Natl. Acad. Sci. USA*

WO 9855132
PCT/US98/10104
treatment of a disease in a mammal, particularly a cow, pig, sheep, mouse or human, and includes:

(a) preventing prion disease or symptoms from occurring in a subject which may be predisposed to the disease or symptom or infected with prion particles but has not yet been diagnosed as having a prion disease which can include the use of gene therapy;

(b) inhibiting prion disease symptoms, i.e., arresting the development of prion disease; or
(c) relieving a prion disease symptom, i.e., causing regression of prion disease or prion disease symptoms.

The term "isolated" shall mean separated away from its natural environment. An isolated protein is not necessarily separated away from all materials it is normally present with and may remain glycosylated.

The term "corresponding position" means the position of an amino acid in a peptide or the position of a codon in a nucleotide sequence corresponds to the same position in the sequence of a different species. The amino acid sequence of PPMF also has corresponding positions from one species to another and corresponding positions for four different positions on the discontinuous epitope of PrP^{Sc} (for five different proteins) are shown in Table 2.

The term "TVB" refers to a mouse strain commonly used in the production of transgenic mice. For purposes of this invention it should be noted that the mouse prion protein (PrP) gene is intact and mouse PrP is therefore expressed at normal levels.

The term "Prnp-%" or "Prnp-Abi" refers to a transgenic animal which has its PrP gene ablated with the "%"; indicating that both alleles are ablated whereas 0/+ indicates only one is ablated. Specifically, the animal being referred to is generally a transgenic mouse which has its PrP gene ablated i.e., a PrP knockout mouse. In that the PrP gene is disrupted no mouse PrP protein is expressed.

The term "sporadic CJD" abbreviated as "sCJD" refers to the most common manifestation of Creutzfeldt-Jakob Disease (CJD). This disease occurs spontaneously in individuals with a mean age of approximately 60 at a rate of 1 per million individuals across the earth.

The term "iatrogenic CJD" abbreviated as "iCJD" refers to disease resulting from accidental infection of people with human prions. The most noted example of such is the

89:9097-9101 (1992) which is incorporated herein by reference to disclose and describe such sequences. The sequences of PrP genes varies from species to species and the differences between a human PrP gene and that of a mouse, cow and sheep are shown respectively in Figures 3, 4 and 5.

5 The terms "standardized prion preparation," "prion preparation," "preparation" and the like are used interchangeably herein to describe a composition containing prions which composition is obtained from brain tissue of mammals which contain substantially the same genetic material as relates to PrP proteins, e.g., brain tissue from a set of mammals which exhibit signs of prion disease which mammals may comprise any of (1) a PrP chimeric transgene; (2) have an ablated endogenous PrP gene; (3) have a high copy number of PrP genes from a genetically diverse species; or (4) are hybrids with an ablated endogenous PrP gene and a PrP gene from a genetically diverse species. The mammals from which standardized prion preparations are obtained exhibit clinical signs of CNS dysfunction as a result of inoculation with prions and/or due to developing the disease due to their genetically modified make up, e.g., high copy number of PrP genes.

15 The term "artificial PrP gene" is used herein to describe a PrP gene which has one or more, preferably two more native codons replaced with different codons wherein the replaced codons are positioned so that, when expressed, form part of the PrP/PPMF binding epitope. The replacement codons preferably express an amino acid different from the native, e.g., basic to acidic, polar to non-polar.

20 The term "chimeric PrP gene" describes recombinantly constructed genes which when included in the genome of a host animal (e.g., a mouse) will render the mammal susceptible to infection from prions which naturally only infect a genetically diverse test mammal, e.g., human, bovine or ovine. In general, an artificial gene will include the codon sequence of the PrP gene of the mammal being genetically altered with one or more (but not all, and generally less than 40) codons of the natural sequence being replaced with a different codon - preferably a corresponding codon of a genetically diverse mammal (such as a human). The genetically altered mammal being used to assay samples for prions which only infect the genetically diverse mammal. Examples of artificial genes are mouse PrP genes encoding the sequence as shown in Figures 3, 4 and 5 with one or more different replacement codons selected from the codons shown in these Figures for humans, cows and sheep replacing mouse codons at the same position, with the proviso that not all the mouse

codons are replaced with differing human, cow or sheep codons. Artificial PrP genes of the invention can include not only codons of genetically diverse animals but may include codons and codon sequences associated with genetic prion diseases such as CJD and codons and sequences not associated with any native PrP gene but which, when inserted into an animal, render the animal susceptible to infection with prions which would normally only infect a genetically diverse animal. In one specific example the chimeric gene is comprised of the starting and terminating sequence (i.e., - and C-terminal codons) of a PrP gene of a mammal of a host species (e.g., a mouse) and also containing a nucleotide sequence of a corresponding portion of a PrP gene of a second mammal of a second species (e.g., a human). A chimeric gene will, when inserted into the genome of a mammal of the host species, render the mammal susceptible to infection with prions which normally infect only mammals of the second species. A useful chimeric gene is MHi2M which contains the starting and terminating sequence of a mouse PrP gene and a nonterminal sequence region which is replaced with a corresponding human sequence which differs from a mouse PrP gene in a manner such that the protein expressed thereby differs at nine residues.

15 The term "genetic material related to prions" is intended to cover any genetic material which effects the ability of an animal to become infected with prions. Thus, the term encompasses any "PPMF gene," "PrP gene," "artificial PrP gene," "chimeric PrP gene" or "ablated PrP gene" which terms are defined herein as well as mutations and modifications of such which effect the ability of an animal to become infected with prions. Standardized prion preparations are produced using animals which all have substantially the same genetic material related to prion so that all of the animals will become infected with the same type of prions and will exhibit signs of infection at about the same time.

20 The terms "host animal" and "host mammal" are used to describe animals which will have their genome genetically and artificially manipulated so as to include genetic material which is not naturally present within the animal. For example, host animals include mice, hamsters and rats which have their endogenous PrP gene altered by the insertion of an artificial gene or by the insertion of a native PrP gene of a genetically diverse test animal.

30 The terms "test animal" and "test mammal" are used to describe the animal which is genetically diverse from the host animal in terms of differences between the PrP gene of the host animal and the PrP gene of the test animal. The test animal may be any animal for which one wishes to run an assay test to determine whether a given sample contains prions to

which the test animal would generally be susceptible to infection. For example, the test animal may be a human, cow, sheep, pig, horse, cat, dog or chicken, and one may wish to determine whether a particular sample includes prions which would normally only infect the test animal. This is done by including PrP gene sequences of the test animal into the host animal, administering PPMF and inoculating the host animal with prions which would normally only infect the test animal.

The terms "genetically diverse animal" and "genetically diverse mammal" are used to describe an animal which includes a native PrP codon sequence of the host animal which differs from the genetically diverse test animal by 17 or more codons, preferably 20 or more codons, and most preferably 28-40 codons. Thus, a mouse PrP gene is genetically diverse with respect to the PrP gene of a human, cow or sheep, but is not genetically diverse with respect to the PrP gene of a hamster. In general, prions of a given animal will not infect a genetically diverse animal and PPMF of a given animal will not bind to PrP^C of a genetically diverse animal.

The terms "ablated prion protein gene," "disrupted PrP gene," "ablated PrP gene," "PrP⁰," and the like are used interchangeably herein to mean an endogenous prion protein gene which has been altered (e.g., add and/or remove nucleotides) in a manner so as to render the gene inoperative. Examples of nonfunctional PrP genes and methods of making such are disclosed in Blümler, H., et al "Normal development of mice lacking the neuronal cell-surface PrP protein" *Nature* 356, 577-582 (1992) which is incorporated herein by reference. Both alleles of the genes are disrupted.

The terms "hybrid animal," "transgenic hybrid animal" and the like are used interchangeably herein to mean an animal obtained from the cross-breeding of a first animal having an ablated endogenous PrP gene with a second animal which includes either (1) a chimeric gene or artificial PrP gene or (2) a PrP gene from a genetically diverse animal. For example a hybrid mouse is obtained by cross-breeding a mouse with an ablated mouse PrP gene with a mouse containing (1) human PrP genes (which may be present in high copy numbers) or (2) chimeric genes. The term hybrid includes any offspring of a hybrid including inbred offspring of two hybrids provided the resulting offspring is susceptible to infection with prions with normal infect only a genetically diverse species.

The terms "susceptible to infection" and "susceptible to infection by prions" and the like are used interchangeably herein to describe a transgenic or hybrid test animal which

develops a prion disease if inoculated with prions which would normally only infect a genetically diverse test animal. The terms are used to describe a transgenic or hybrid animal such as a transgenic mouse Tg(MHu2M) which, without the chimeric PrP gene, would not be susceptible to infection with a human prion (less than 20% chance of infection) but with the chimeric gene is susceptible to infection with human prions (80% to 100% chance of infection).

The terms "resistant to infection", "resistant to infection with prions" and the like mean the animal includes a PrP gene which renders the animal resistant to prion disease when inoculated with an amount and type of prion which would be expected to cause prion disease in the animal. The resistant animals PrP gene includes non-native codons which express amino acids different from those of the native PrP gene which effect the PrP^{Sc}/PPMF binding site.

The term "incubation time" shall mean the time from inoculation of an animal with a prion until the time when the animal first develops detectable symptoms of disease resulting from the infection. A reduced incubation time is six months or less, preferable about 100 days \pm 25 days or less, more preferably about 30 days \pm 10 days or less.

ABBREVIATIONS USED HEREIN INCLUDE:

- BSE for bovine spongiform encephalopathy;
- CJD for Creutzfeldt-Jakob Disease;
- CNS for central nervous system;
- FFI for fatal familial insomnia;
- FVB for a standard inbred strain of mice often used in the production of transgenic mice since eggs of FVB mice are relatively large and tolerate microinjection of exogenous DNA relatively well;
- GSS for Gerstmann-Strassler-Scheinker Disease;
- Hu for human;
- HuPPMF for human Prion Protein Modulator Factor;
- HuPrP for a human PrP protein;
- MHu2M for a chimeric mouse/human PrP gene wherein a region of the mouse PrP gene is replaced by a corresponding human sequence which differs from mouse PrP at 9 codons;

MHu2MPp^{Sc} for the scrapie isoform of the chimeric human/mouse PrP gene;
Mo for mouse;

MoPPMF for mouse Prion Protein Modulator Factor;

Mo PrP for a mouse PrP protein;

MoPrP^{Sc} for the scrapie isoform of the mouse PrP protein;

PPMF for Prion Protein Modulator factor in general, i.e., that protein as in any species;

Prp^{Sc} for ablation of both alleles of an endogenous PrP protein gene, e.g., the Mo PrP gene;

Prp^{Sc} for the CJD isoform of a PrP gene;

Prp^{Sc} for the scrapie isoform of the PrP protein;

ScN2a for persistently infected scrapie mouse neuroblastoma cells also expressing (MHu2)PrP^{Sc};

SHa for a Syrian hamster;

SHa PrP for a Syrian hamster PrP protein;

Tg for transgenic;

Tg(BovPrP) for transgenic mice containing the complete cow PrP gene;

Tg(HuPrP) for transgenic mice containing the complete human PrP gene;

Tg(HuPrP)/Prp^{Sc} for a hybrid mouse obtained by crossing a mouse with a human PrP protein gene (HuPrP) with a mouse with both alleles of the endogenous PrP protein gene disrupted;

Tg(MHu2M) mice are transgenic mice of the invention which include the chimeric MHu2M gene;

Tg(MHu2M)/Prp^{Sc} for a hybrid mouse obtained by crossing a mouse with a chimeric PrP protein gene (MHu2M) with a mouse with both alleles of the endogenous PrP protein gene disrupted;

Tg(SHa PrP) for a transgenic mouse containing the PrP gene of a Syrian hamster;

Tg(SHa PrP^{Sc}81/Prp^{Sc}) for a particular line (81) of transgenic mice expressing SHA PrP, +/0 indicates heterozygous; and

Tg(SHa PrP) for transgenic mice containing the complete sheep PrP gene.

GENERAL ASPECTS OF THE INVENTION

An understanding of PrP proteins and their characteristics provides background information to understand the function and characteristics of PPMF and equivalent pharmacophores. When the PrP gene is normally expressed it produces PrP^C which is the protein in its "cellular" non-disease form. In order to develop a prion disease PrP^C must be converted to PrP^{Sc} which is the "scrapie" or disease causing form of the protein. Both PrP^C and PrP^{Sc} have the same amino acid sequence but a different 3-dimensional conformation.

PrP^{Sc} is referred to as a prion or infectious protein because when a mammal such as a mouse is inoculated with mouse prions (i.e., mouse PrP^{Sc}) the mouse will develop prion disease. However, prions are species specific meaning that if a mouse is inoculated with human or cow prions the mouse will not develop disease.

To begin characterization of PPMF it was noted that a transgenic mouse containing a human PrP gene would not get sick when inoculated with human prions. However, if the mouse PrP gene is ablated and the mouse genome includes the human PrP gene the mouse will get sick when inoculated with human prions. It was also noted that when a transgenic mouse was created with a chimeric (part mouse and part human) PrP gene the mouse would get sick when inoculated with human prions.

The above information was combined with results obtained with murine ScN2a cells showing that human COOH-terminal residues prevented chimeric human/mouse PrP^C from being converted to the PrP^{Sc} form.

Next it was found that by making particular substitutions in a PrP gene it is possible to prevent the conversion of PrP^C to PrP^{Sc}. Specifically, it was found that by substituting human residues I215 or E2K219 for mouse V or Q respectively (at a corresponding position) the conversion of mouse PrP^C into PrP^{Sc} was prevented. Further, the substitution of R for Q171 in a mouse PrP gene prevents the formation of MoPrP^{Sc}. Substitutions at these relatively remote positions indicated that the PrP^C protein was folded such that these positions were moved together. This was confirmed by producing an NMR image showing residues 170 and 171 in a loop adjacent to residues 214 and 218 on the PrP^C protein (see Figure 2).

In order for PrP^C to convert to PrP^{Sc} the PrP^C must bind PPMF. Residues in the vicinity of 171 and 215 form a discontinuous epitope to which PPMF binds. If that binding is prevented (or occurs in a manner such that the PPMF is not released) then PrP^C is not

converted to PrP^{Sc}. By knowing the general area of the discontinuous binding epitope for PPMF on PrP^C it is possible to replace codons (and thus amino acid residues) in that area and thereby effect PPMF/PrP^C binding. However, knowledge of the linear sequences of amino acids in the area of the discontinuous epitope does not completely define the epitope.

Figure 2 shows the area of the discontinuous epitope on PrP^C for binding of PPMF which area shows that the PrP^C epitope is not a linear binding site. Specifically, Figure 2 shows that the 215 and Gln 219 are on the outwardly facing surface of the discontinuous epitope as is Gln 172 - these positions are for hamster PrP^C. The figure shows that due to the loops formed, the residues immediately before and after 215 and 219 are facing away from the surface presented by the PrP^C epitope. Thus, it is not a linear sequence of amino acids which define the binding epitope surface but rather a surface defined in 3-dimensional space by the distances and angles shown in Figure 4. The negative image of that surface defines the surface of PPMF to which the PrP^C epitope binds.

15 IDENTIFYING PHARMACOPHORES

Figures 12-15 are schematic flow diagrams of steps involved in identifying compounds which inhibit PrP^{Sc} formation. The process begins with the identification of the predicted PPMF binding site of the PrP^C gene of a particular species. Once this is accomplished, a large chemical directory which may contain any number of compounds is searched for compounds that may interact with the species-specific PPMF binding site of the PrP^C. For example, the initial group of identified compounds may contain tens or hundreds of thousands or more of different non-peptide organic compounds. A different or supplemental group may contain millions of different peptides which could be produced synthetically in chemical reactions or via bacteria or phage. Large peptide libraries and methods of making such are disclosed in U.S. Patents 5,266,684, issued November 30, 1993, and 5,420,246, issued May 30, 1995, which are incorporated herein by reference. Libraries of non-peptide organic molecules are disclosed in PCT publication WO 96/40202, published December 19, 1996, incorporated herein by reference.

The initial library of compounds is screened via computer generated modeling, e.g., computer models of the compounds are matched against a computer model of the PPMF binding site on PrP^C (see Figure 17) to find molecules which mimic the spatial orientation

and basic polymorphism of PPMF. This screening should substantially reduce the number of candidate molecules relative to the initial group.

The screened group is then subjected to further screening visually using a suitable computer program which makes viewable images of the molecules. The resulting candidate molecules are then actually tested for their ability to inhibit PrP^{Sc} formation. This testing is described in Example 20. As a result of the assays carried out, six compounds were found to inhibit PrP^{Sc} formation. The six compounds are identified in the Compound Table which shows that four of the six compounds are cytotoxic to the ScN2a cells while two compounds are not cytotoxic and inhibit PrP^{Sc} formation. Figure 16 is a computer generated diagram showing the six compounds which inhibit PrP^{Sc} formation and specifically showing how and where each compound binds to the PPMF binding site on PrP^C (90-231).

CHARACTERIZATION OF PPMF

Prion Protein Modulator Factor (PPMF) is a species specific protein which binds to PrP^C and facilitates a conformational change from PrP^C to PrP^{Sc}. PPMF is further characterized by not binding to PrP^{Sc} and acting as rate-limiting component in the conversion of PrP^C to PrP^{Sc}. Specific species of PPMF are encoded by specific nucleotide sequences derived from a mammal selected from the group consisting of human, cow, sheep, mouse and hamster. PPMF is further characterized by diminishing the activation energy barrier between PrP^C and PrP^{Sc} thereby facilitating the formation of PrP^{Sc}. In addition to these generalized characteristics, PPMF has the following specific identifying characteristics. Corresponding positions between the residues for different PrP proteins are provided in Figures 3, 4 and 5. Corresponding positions for five different PrP proteins are shown in Table 2. PPMF is not affected by a change in residue at position 216, 220, 221 or 222 of human PrP^C or a corresponding position in a different species. As shown in Figure 2, the position next to a position which affects binding is likely to be facing away from the epitope surface. The binding of PPMF to PrP^C is affected by residues at position 215 and 219 in hamster PrP^C and corresponding positions in the PrP^C protein of different species. Binding of PPMF to PrP^C is affected by the residue at position 172 in sheep PrP^C and the corresponding residue in the PrP^C protein of different species. This would generally not be expected as the position 171 is some distance from the 214-219 region. However, the folding of PrP^C proteins as per Figure 2 explains this. Finally, PPMF binds to a

discontinuous epitope on PrP^C which epitope comprises residues 172, 215 and 219 in the human PrP protein and corresponding residues in the PrP protein of different species.

USE AND OPERATION OF PPMF

5 In order for a mammal to develop a prion disease, PrP^C must be converted to PrP^{Sc}, i.e., prions must be formed. In order for prions to be formed, three compounds must be present which are PrP^C, PrP^{Sc}, and PPMF. Because PPMF is a rate-limiting compound in the formation of prions, (if not recycled) an animal which is infected with prions (i.e. PrP^{Sc}) may develop symptoms of prion disease very slowly. This is not desirable when the animal is being used as a test animal in order to determine if the prions are present within a sample. Thus, the administration of PPMF to such test animals can greatly reduce the amount of time necessary for the formation of prions and thereby reduce the amount of time necessary to pass before the observation of the first symptoms of prion disease.

15 Figure 1 schematically shows how PPMF is involved in the conversion of PrP^C to PrP^{Sc}. Specifically, PrP^C is represented by the circle 1 and is present in a non-diseased normal animal as is PPMF represented by the C-shaped molecule 2. When PPMF binds to PrP^C it allows for the formation of a complex which includes PrP^{Sc} represented by the squares 3. After the complex is formed the PrP^C is converted to PrP^{Sc} and the PPMF is released as it does not bind PrP^{Sc}.

20 Based on both an understanding of how PrP^C is converted to PrP^{Sc} as shown in Figure 1 and the 3-dimensional conformation of the PrP^C binding epitope as shown in Figure 4 it is possible to construct a variety of therapeutic compounds which allow for the treatment of prion disease. The specific amino acid positions and coordinates of Figure 4 are provided with respect to hamster PrP^C. However, using Table 2 it is possible to substitute the amino acids shown for corresponding positions for human, mouse, cow and sheep and the chimeric PrP protein. With this information it is possible to generate a variety of pharmacophores which mimic the epitope surface of PrP^C. Such pharmacophores will bind to the binding surface of PPMF and as such can be used to assay for the presence of PPMF, determine its concentration within a sample, and, more importantly, be used as a therapeutic for the treatment of prion disease.

30 By knowing the amino acids positioned on the surface of the PrP^C epitope and knowing the precise 3-dimensional position of the residues as shown within Figure 4, the 3-

dimensional surface of the binding surface of PPMF can be generated. The binding surface of PPMF to PrP^C is, of course, the negative image of the epitope binding surface of PrP^C. By knowing the negative image of the epitope surface of PrP^C (i.e., the 3-dimensional coordinates of the binding surface of PPMF) it is possible to generate PPMF pharmacophores. The PPMF pharmacophores have a variety of specific uses. Firstly, the pharmacophores of PPMF can be bound to the surface and be used to assay for the presence of PrP^C in that they bind to the PrP^C epitope. Secondly, when the pharmacophores are designed so as to allow for the conversion of PrP^C to PrP^{Sc} they can be administered to two animals such as mice which are being inoculated with a sample in order to test for the presence of prions within the sample. By administering the pharmacophores, the time needed to develop symptoms of prion disease will be greatly reduced provided the sample used for inoculation actually does contain PrP^{Sc} (i.e., contains prions). Perhaps most importantly, the pharmacophores can be designed to bind tightly to the PrP^C epitope so that they are not released from the epitope and do not result in the conversion of PrP^C to PrP^{Sc} -- and such pharmacophores would be valuable therapeutics for the treatment of prion disease.

5 PPMF is useful in enhancing the speed and sensitivity of an assay which uses a transgenic animal to detect the presence of PrP^{Sc} (i.e., detect prions) in a sample. Specifically, PPMF is administered to transgenic animals which have been inoculated with a sample which may contain prions. The transgenic animals are genetically designed so that they are susceptible to infection by prions which would normally only infect a genetically diverse species. The animals are observed to determine if they develop symptoms of prion infection which if observed allows the tester to deduce that the sample includes prions. The administration of PPMF enhances the rate of disease development thereby shortening assay time.

15 PPMF is useful in enhancing the speed and sensitivity of an assay which uses a transgenic animal to detect the presence of PrP^{Sc} (i.e., detect prions) in a sample. Specifically, PPMF is administered to transgenic animals which have been inoculated with a sample which may contain prions. The transgenic animals are genetically designed so that they are susceptible to infection by prions which would normally only infect a genetically diverse species. The animals are observed to determine if they develop symptoms of prion infection which if observed allows the tester to deduce that the sample includes prions. The administration of PPMF enhances the rate of disease development thereby shortening assay time.

25

ASSAYS

Assays for detecting the presence of PrP^{Sc} in a sample can be facilitated by the use of PPMF, pharmacophores thereof and/or antibodies which bind to PrP^C at its PrP^C/PPMF binding site. The sample suspected of containing PrP^{Sc} is obtained and prepared as needed. Then PrP^C and PPMF are added which will facilitate the formation of PrP^C/PPMF/PrP^{Sc} complexes if PrP^{Sc} is present in the sample. Once the complex is formed the PrP^C is converted to PrP^{Sc} and the complex breaks up allowing the formation of two more

complexes. The reaction proceeds geometrically increasing the number of Prp²⁶ molecules in the sample i.e., 2, 4, 8, 16, etc., for each Prp²⁶ molecule present. After the reaction is allowed to sufficiently proceed it can be tested for the presence of Prp²⁶ using any desired assay such as an assay disclosed in U.S. Patent Application Serial No. 08/804,536, filed February 21, 1997, which is incorporated herein by reference.

The assay can be used to test for therapeutics. Prp²⁶ and PPMF are added to a known concentration of Prp²⁶ and a compound to be tested is added. If the compound prevents the formation of Prp²⁶ the compound is a potential therapeutic for the treatment of prion diseases. The ability to assay for increasing in the concentration of Prp²⁶ can be determined using an assay and equation disclosed and described in U.S. Patent Application Serial No. 08/804,536, filed February 21, 1997.

ASSIGNMENT OF THE BINDING SITE OF PPMF ONTO THE Prp²⁶ STRUCTURE BY NMR

Based on experiments carried out Mo residues 214 and 218 (corresponding to SHa 215 and 219) were assigned to the NMR structure of recombinant SHa Prp 90-231. These residues form a binding pocket-like structure facing to a plasma membrane to which Prp²⁶ is anchored via GPI anchor (Figure 2). In this pocket, SHa residues 215 and 219 (214 and 218 in Mo) are located on its surface, whereas SHa residue 217 (216 in Mo) is buried and not exposed as our biochemical data indicated. Interestingly, SHa residues 165-173 (164-173 in Mo) on a loop are also participating in this binding pocket-like formation. Gln/Arg polymorphism at sheep residue 171 (167 Mo) has been reported to determine its susceptibility to sheep natural scrapie as Hu polymorphism at residue 219 (218 in Mo) does. Based on the computer assignment, Lys substitution at Hu residue 219 showed an equivalent effect to Gln to Arg substitution at sheep residue 171 on this pocket formation.

PPMF is further characterized by the information provided in Table 1 below.

TABLE 1

Prp ²⁶	Prp ²⁶ formation	MHM2 Prp ²⁶ formation when co-transfected	Binding affinity to PPMF
SHa	++	++	-
MH3	++	++	-
MH2M	+	+	-
MHM2	-	-	-
Mo	-	-	-
MH3HuA	(-)	-	(-)
MH2HuA	(-)	-	(-)
MHMHuA	(-)	-	(-)
M3HuA	(-)	-	(-)
MHMHuA3	++	-	++
MHMK218	(-)	++	+
MHM1218	+	-	+
MHMA218	++	++	+
MHMW218	+	++	+
MHMF218	(-)	--	--
MHMR216	(-)	++	-
MHMFHu216	(-)	-	++
MHMHuA1	+	++	+
MHMK214	(-)	-	(-)
MHME214	(-)	++	+
MHMA214	-	-	-
MHMW214	(-)	++	+
MHMF214	(-)	-	(-)

The data presented here in combination with the previous notions obtained by Tg studies allows the following conclusions regarding Prp²⁶ formation. Mo Prp²⁶ does not inhibit the MHM2 Prp²⁶ formation when co-transfected because PPMF can be released from

the Mo PrP^C after the PrP^C is converted into PrP^{Sc}. PPMF is not able to bind to PrP^{Sc} efficiently. In contrast, SHa or MH3 PrP^C remain in the PrP^C isoform and capture PPMF in PrP^C-PrP^{Sc} complexes. This results in the depletion of uncomplexed PPMF. In the case of HuA derivatives, PPMF remains free and unbound since the unconverted PrP^C/PrP^{Sc} complexes have no binding site for PPMF regardless of the compatibility at its central region. This shows that PPMF is a less abundant protein and is a rate-limiting substrate if not recycled. This shows that the levels of PPMF must be <50% of the levels of endogenous PrP^C to convert PrP^C to PrP^{Sc} and result in disease. Overexpression of SHa or Mo PrP^C increased the rate of PrP^{Sc} formation indicating that PPMF is not rate-limiting and is recycled.

A Hu polymorphism at residue 219 (corresponding to Mo 218) has been reported in the Japanese population. About 12% of the people have the Lys allele instead of the Glu/Glu allele, but so far none of autopsy-proven Japanese patients with CJD possessed the Lys allele at Hu residue 219. The absence of Lys 219 allele in Japanese patients with CJD is due to an aberrant binding affinity of the PrP^C to PPMF. Even if PrP^C carrying Lys allele is unable to convert into PrP^{Sc}, one might think that the counterpart of wt-PrP^C could be a substrate to convert into PrP^{Sc}, which means allele frequency must be half (6%) in the Japanese CJD population. However, the Examples provided here also revealed that the Lys allele exhibited even tighter binding to PPMF so that the free uncomplexed PPMF is depleted. This explains why the counterpart of wt-PrP^C does not convert into PrP^{Sc}. Another sheep Gln/Arg polymorphism at sheep residue 171 also determines its susceptibility to sheep natural scrapie. Knowing that this site is also participating in the binding pocket-like formation as shown in the NMR structure shows that susceptibility to the sheep natural scrapie is governed by PrP^C-PPMF interaction at the residue.

One urgent application from the current findings is for the therapeutic approach in prion disease. PPMF is present in such small amounts (<50% of endogenous PrP^C) that it is a rate-limiting factor in the formation of PrP^{Sc} if not recycled. Modifying the PPMF binding site in PrP^C efficiently blocks conversion of its own or co-expressed wt-PrP^C into PrP^{Sc}. Thus, it is clear the binding site for PPMF primarily governs the inhibition in PrP^{Sc} formation over a segment of PrP^C including the central region between residues to 96-167 mouse PrP^C. Molecules which tightly bind PPMF have therapeutic applications in the

treatment of prion disease. Further, both human and sheep polymorphisms at the binding pocket for PPMF regulate susceptibility to prion disease.

CORRESPONDING AMINO ACID POSITIONS

- 5 Mammals possess a PrP gene which expresses a PrP protein in the "cellular" form, i.e., PrP^C. With respect to some mammals such as mice and hamsters, the PrP gene and resulting PrP protein are genetically similar. However, with respect to other species such as mice and humans, the PrP gene is genetically diverse.
- 10 When a normal animal is inoculated with prions taken from the same or a genetically similar species, the animal will become sick with prion disease. However, when an animal is inoculated with prions from a genetically diverse species the animal does not become sick with prions. This species barrier relates to the species specificity of PPMF and the binding epitope on PrP^C. Although much of the information provided here such as that shown within Figures 7 and 9 is specific to hamster PrP^C the information is generally applicable to other species by utilizing the information provided below in Table 2 which shows the
- 15 corresponding amino acid positions for the four critical residues for human, mice, hamster, cow, sheep and the chimeric human mouse PrP gene. Corresponding positions are shown in Figure 10 along with their relative positions on the PrP protein structure.

TABLE 2

MUTATIONS IN ENTROPE TAGGED MINE PRP INHIBIT PRP ⁺						
FORMATION IN SC2A CELLS						
Mo Codon Number	PrP Residue: Mouse	Human	Syr Hamster	Sheep	Mutant MHM2	Type of inhibition of PrP ⁺ Formation
167	Q	E	Q	Q/R		
170	N				S	None
171	Q	Q	Q	Q	R	2
214	V	I	T	I	I	2
					K	1
					E	2
					A	1
					W	2
					P	1
215	T	T	T	T	Q	None
216	Q	Q	Q	Q	R	2
218	Q	E/K	Q	Q	E	1
					K	2
					I	1
					A	2
					W	2
					P	1
					F	1
					R	2
					H	2
219	K	R	K	R	R	None
221	S	S	S	S	A	None
222	Q	Q	Q	Q	K	None

TABLE 2 (Continued)

MECHANISMS OF INHIBITION OF PrP ⁺ FORMATION			
Type of Inhibition	Example	Positive Mechanism	Relative affinity for protein X*
5	1	HuPrP ^C (E219) binding to Mo protein X inhibited by MoPrP ^C	Competitive
			Low
10	2	HuPrP ^C (K219) prevents MoPrP ^C binding to protein X	Non-competitive
			High
	3	ShPrP ^C binds to protein X and is not released by MoPrP ^C	Non-competitive
			Similar
10		*affinity relative to that of MoPrP ^C for Mo protein X.	

PRION RESISTANT ANIMALS

Animals resistant to prion disease could be created by (1) ablating the PrP gene; (2) making substitutions in the PrP gene at critical points within the PrP/PPMF binding site or (3) examining the PrP gene of animals and breeding those animals which are found to include codons which encode amino acids at the PrP epitope site which would render the animal resistant to prion disease.

Although ablating the PrP gene might be the simplest route to obtain a prion resistant animal, such has been accomplished only in mice -- although others have suggested that the PrP gene could be ablated in other animals (see PCT publication WO93/10227 published May 27, 1993).

PrP genes have been extracted and isolated. Using the site directed recombination (homologous recombination methods), the codons at particular positions can be replaced to create artificial genes of the invention which genes, when operably inserted in the genome of an animal, render the animal resistant to prion infection. Using the information within Table 2 and Figure 4, it is possible to carry out codon substitutions at all or any of the four sites within the PrP epitope which would affect binding of the PrP protein to PPMF. The effect of a number of substitutions are shown in Table 2. After creating the desired recombinant PrP gene, the gene could be injected into a fertilized egg and implanted in a pseudo-pregnant female and allowed to grow to term. If the implanted recombinant PrP gene replaced the native PrP gene, a transgenic animal results which is resistant to prion disease.

- (b) the binding is species specific, i.e. PPMF of a given species binds to PrP^C of that species only on a species which is genetically similar in terms of its PrP^C proteins;
- (c) does not bind to PrP^C and as such is released from a PrP^C molecule after it is converted to PrP^{Sc};
- (d) acts as a rate limiting component in the conversion of PrP^C to PrP^{Sc};
- (e) binding of PPMF to PrP^C is not effected by residues 216, 220, 221 and 222 in human PrP^C protein and the corresponding residues in other species;
- (f) PPMF allows for the formation of a PrP^CPrP^{Sc} complex where the PPMF holds the complex together until PrP^C is converted to PrP^{Sc};
- (g) PPMF is released from the PrP^CPrP^{Sc} complex once PrP^C is converted to PrP^{Sc} as PPMF does not bind to PrP^{Sc};
- (h) PPMF decreases the activation energy barrier between PrP^C and PrP^{Sc};
- (i) binding of PPMF to PrP^C is effected by residues at position 215 and 219 in hamster PrP^C and corresponding positions in the PrP^C protein of other species;
- (j) binding of PPMF to PrP^C is effected by a residue at position 172 in S14a PrP^C and a corresponding position in a different species;
- (k) PPMF binds to a discontinuous epitope on PrP^C which epitope comprises residues 172, 215, 219 in human PrP^C and the residues at corresponding positions at the PrP^C protein of other species;
- (l) preferred forms of PPMF are encoded by a nucleotide sequence of a species selected from the group consisting of human, cow, sheep, mouse and hamster.

PHARMACOPHORES

- The shape that PrP^C adopts at its binding epitope when bound to the biologically active target molecule PPMF, the biological shape, is an essential component of its biological activity. This shape, and any specific interactions such as hydrogen bonds, can be exploited to derive predictive models used in rational drug design. These can be used to optimize lead compounds, designed *de novo* compounds, and search databases of existing compounds for novel structures possessing the desired biological activity. In order to aid in the discovery of useful pharmacophores for the PrP^C binding epitope, these models must make useful predictions, relate chemical structures to activity, and confidently extrapolate to chemical classes beyond those used for model derivation.

- There are a number of mutations and polymorphisms existing with respect to the PrP gene of a different species. A number of the mutations and polymorphisms are listed in the "Mutation Table" provided below. It is believed that additional mutations and polymorphisms exist in all species within the PrP gene. However, until now one could not determine which sequences might provide for resistance to prion disease. However, by using the information within Figure 10 along with the information known with respect to PrP gene sequences, it is possible to screen animals with respect to the sequence of their PrP gene. The screening is carried out in order to determine mutations existing at the sites which match the critical sites indicated in Figure 10 and/or other sites which form part of the PrP^C binding epitope, i.e., the epitope on PrP^C which binds to PPMF. Animals with a PrP gene which is heterozygous at a particular point could be bred with other animals which are heterozygous at that point in order to produce offspring which include those with a homozygous PrP gene of the type desired.

- Whether screening or breeding animals or producing recombinant PrP genes, certain factors should be considered. Specifically, it is most desirable to breed for an animal or produce a PrP gene wherein more than 1 of the critical sites as per Figure 10 is substituted with a new codon which will produce a new amino acid. Secondly, the substitution at the critical site should be with an amino acid which is biochemically quite different from the amino acid at that position which is known to render the animal susceptible to prion infection. Thus, if a basic and/or polar amino acid is present at the critical site that site could be replaced with an acidic and/or nonpolar amino acid. With these criteria in mind some trial and error would be required. However, by knowing the critical positions per Table 2 and Figures 7 and 15 and knowing that substitutions should be made to make the amino acid different, the desired results can be obtained. Acidic amino acids should be substituted with basic amino acids and vice versa. Polar amino acids should be substituted with nonpolar amino acids and vice versa.

PRION PROTEIN MODULATOR FACTOR (PPMF)

- Based on experiments and results described here and deductions made therefrom, the protein PPMF is characterized as follows:
- (a) binds to PrP^C and facilitates a conformational change from PrP^C to PrP^{Sc};

Pharmacophore models (e.g., BioCAD incorporated herein by reference) model activity in terms of the positions of a small number of atoms of particular functional groups. This overcomes many of the problems of traditional QSAR models. U.S. Patent 5,025,388 to Kramer *et al.* provides for comparative molecular field analysis (COMFA incorporated herein by reference) methodology. In accordance with this methodology the 3-dimensional structure for each molecule is placed within a 3-dimensional lattice and a probe atom is chosen, placed successively at each lattice intersection and the steric and electrostatic interaction energies between the probe atom and the molecule are calculated for all lattice intersections. The energies are listed in a 3-dimensional-QSAR table. A field fit procedure is applied by choosing the molecule with the greatest biological activity as the reference in conforming the remaining molecules to it.

The methodology disclosed within U.S. Patent 5,526,281 (incorporated herein by reference in its entirety) is particularly useful for the generation of pharmacophores using the information provided herein and in particular the 3-dimensional coordinates shown within Figure 4 which can be applied to the PrP^C protein of other molecules using the information of Figure 10. This methodology is particularly useful for the generation of pharmacophores of the present invention because in many binding interactions between molecules, not all the characteristics of the molecule considered are of equal importance. As shown within Figure 2, many of the amino acids within the PrP^C protein are of little importance with respect to the surface of the binding epitope presented for binding PPMF. The approach disclosed within 5,526,281 allows the user to focus on the salient features of the molecule.

The first step in the '281 method for generating pharmacophores involves the selection of a pose. A pose of a molecule is defined by its confirmation (internal torsional angles of the rotatable bonds) and orientation (the rigid rotations and translations). This mathematically defines the pose of a molecule and in connection with the present invention is shown within Figure 4. Figure 4 shows the coordinates for a pose for the binding epitope of PrP^C. The negative image of the coordinates as shown in Figure 4 can also be generated and as such represents a pose for the binding epitope of PPMF. The negative image and other possible pharmacophores can be generated using software available such as CatalystTM from BioCad, Foster City, California and, BatchminTM available from Columbia University, New York City, New York (both of which are incorporated herein by reference). These

programs take into consideration various properties including physical and chemical properties, Shape, electrostatic interaction, solvation and biophysical properties.

Other methods for generating pharmacophores of the present invention are disclosed within U.S. Patent 5,307,287 issued April 26, 1994 and U.S. Patent 5,434,796 issued July 18, 1995 (both of which are incorporated herein by reference in their entirety). Comparative molecular field analysis (CoMFA) is an effective computer implemented methodology of 3D-QSAR employing both interactive graphics and statistical techniques for co-relating shapes of molecules with their observed biological properties. For each molecule in a series of known substrates the steric and electrostatic interaction energies with a probe atom are calculated at special coordinates around the molecule. Subsequent analysis of the data table by a partial list square (PLS) cross-validation technique yields a set of coefficients which reflect the relative contribution of the shape elements of the molecular series to differences in biological activities. Display in three dimensions in an interactive graphics environment of the special volumes highly associated with biological activity, and comparison with molecular structures yields an understanding of intermolecular associations. CoMFA will also predict the biological activity of new molecular species such as various pharmacophores of either the PrP^C binding epitope or the surface to which it binds on PPMF.

TEST ANIMAL

Although a variety of different test animals could be used for testing for the presence of prions within a sample, preferred test animals are mice and hamsters, with mice being most preferred in that there exists considerable knowledge on the production of transgenic animals. Other possible host animals include those belonging to a genus selected from *Mus* (e.g. mice), *Rattus* (e.g. rats), *Oryzologus* (e.g. rabbits), and *Mesocricetus* (e.g. hamsters) and *Cavia* (e.g. guinea pigs). In general mammals with a normal full grown adult body weight of less than 1 kg which are easy to breed and maintain can be used. The host PrP gene can be changed to include codons from genetically diverse PrP genes from test animals belonging to a genus selected from *Bos*, *Ovis*, *Sus* and *Homo*. Preferably, a mouse host PrP gene is changed to include codons from a human, cow or sheep PrP gene, with human being most preferred. Humans are preferred because an important object of the invention is to use the animal to test a sample of material to determine if that material has prions which will infect a human and cause a human to develop a CNS disease such as CJD. Preferred

transgenic animals are disclosed in U.S. Patent 5,565,186 issued October 15, 1996 and WO 97/04814 published February 13, 1997 which are incorporated herein by reference to disclose transgenic animals and methods of making and using such.

PCT/US98/10104

The genetic material which makes up the PrP gene is known for a number of different species of animals (see Gabriel et al., *Proc. Natl. Acad. Sci. USA* 89:9097-9101 (1992)). Further, there is considerable homology between the PrP genes in different mammals. Note that the segment of a PrP gene used to create the MHu2M gene of the present invention will result in encoding of protein which shows a difference between the human and a mouse protein of only nine residues. Although there is considerable genetic homology with respect to PrP genes, the differences are significant in some instances. More specifically, due to small differences in the protein encoded by the PrP gene of different mammals, a prion which will infect one mammal (e.g. a human) will not normally infect a different mammal (e.g. a mouse). Due to this "species barrier," it is not generally possible to use normal animals, (i.e., animal which have not had their genetic material related to prions manipulated) such as mice to determine whether a particular sample contains prions which would normally infect a different species of animal such as a human. The present invention solves this problem in a surprising manner.

RELATIONSHIPS - PrP GENES: COPY NUMBERS: GENETIC DIVERSITY

Commercially useful transgenic animals are preferably small and easy to reproduce; thus, host animals such as mice, hamsters, guinea pigs and rats are preferred, with mice being most preferred. In order for the transgenic animals to be useful, it is necessary for the animals to be susceptible to infection with prions which normally infect only genetically diverse test animals, and in particular animals of commercial significance for testing, such as humans, cows, horses, sheep, pigs, cats, dogs and chickens, with humans being most preferred. Further, for the transgenic and hybrid animals to be useful in a practical and commercial sense, it is necessary for the animals to demonstrate symptoms of the disease within a relatively short period after inoculation, and for a very high percentage of the animals to demonstrate symptoms of the disease after inoculation. Both criteria are facilitated by the administration of PPMF.

In producing a transgenic animal having the above-described characteristics, we noted a number of relationships of significance.

-33-

PCT/US98/10104

First, when the entire PrP gene of a test animal (such as a human) is made functional in the host animal (such as a mouse) the resulting transgenic animal (with a low copy number of human PrP genes) is not susceptible to infection with human prions.

Second, we found that infection would occur if the endogenous PrP gene of the host animal is ablated.

Third, when only some of the codons differing between the host and the test animal are switched, the resulting transgenic animal is susceptible to infection with prions which normally only infect the test animal.

Fourth, we noticed that, as the copy number of the artificial gene in the transgenic animal is increased, the incubation time in some cases decreases.

Fifth, we noted that humans with some genetic defects resulting in prion diseases have different genetic defects in their PrP gene and that by matching the defects in any transgenic animal will render that animal more susceptible to infection with prions from the diseased human.

With this knowledge, we deduced that it is possible to produce a transgenic animal wherein all of the PrP gene of the host animal is replaced with the PrP gene of a test animal to obtain a useful transgenic animal which is susceptible to infection with prions which normally only infect the test animal by either (1) administering PPMF and/or (2) substantially increasing the copy number of the test animal's PrP gene in the host animal and preferably also ablating the endogenous PrP gene.

For example, a transgenic mouse which includes the entire PrP gene of a human in a relatively low copy number (e.g. 1 to 4) is not susceptible to infection with human prions (unless the endogenous mouse PrP gene is ablated). However, if the transgenic mouse includes a very high copy number of a human gene (e.g. 30-300 copies), the resulting transgenic animal is susceptible to infection with human prions. Further, when a host animal such as a mouse has only a portion of its PrP gene replaced with a corresponding portion of a test animal such as a human, the resulting transgenic animal is highly susceptible to infection with prions which normally infect only the test animal. This is true even if the chimeric gene is present in the transgenic animal in a relatively low copy number (e.g. 1 to 4 copies) resulting in low expression of MHu2M PrP^C.

Lastly, in order to reduce incubation time, hybrid mice were created by crossing mice with ablated PrP genes with transgenic mice which (1) included a PrP gene from a

-34-

genetically diverse animal e.g., a human or (2) include a chimeric or artificial gene of the present invention. The chimeric gene and/or PrP gene from genetically diverse test animal may be present in high copy number. The copy number can be increased in order to reduce incubation time provided the copy number is not increased so far that the animal becomes spontaneously ill, i.e., become ill without inoculation with prions.

Based on the above, it can be understood that the preferred transgenic animals are (1) animals such as mice which include a chimeric PrP gene, i.e., only a portion, but not all, of their PrP gene replaced with a corresponding portion of the PrP gene of a test animal or (2) animals with an ablated endogenous PrP gene and a PrP gene from another animal such as a human most preferable where that human PrP gene has a genetic defect which results in a prion disease when in a human.

SPECIES BARRIER BROKEN

The transmission of human CJD to apes and monkeys 1.5 - 3 years after intracerebral inoculation provided considerable interest in the causes of neurodegenerative diseases [Gibbs, Jr. et al., *Science* 161:388-389 (1968)]. Humans are not genetically diverse from apes and monkeys which accounts for the cross-species infectivity, although with a long incubation time. While the high cost of caring for nonhuman primates prevented extensive studies of the human prion diseases, the transmissibility of these diseases stimulated studies of the animal prion analogues in rodents [Manuelidis et al., *Proc. Natl. Acad. Sci. USA* 75:3422-3436 (1978); Manuelidis et al., *Proc. Natl. Acad. Sci. USA* 73:223-227 (1976); Tateishi et al., *Ann. Neurol.* 3:581-584 (1979)].

The present disclosure of PPMF allows assays to be performed relatively rapidly in genetically altered mammals such as Tg(MHu2M) mice that are relatively inexpensive to maintain. Endpoint titrations of prions in multiple human body tissues and fluids can be performed and standard curves constructed for more economical incubation time assays. The information derived from such studies of human prions will be useful in the management of CJD patients who are thought to pose some risk to relatives, physicians, nurses and clinical laboratory technicians [Berger et al., *Neurology* 43:205-206 (1993); Ridley et al., *Lancet* 341:641-642 (1993)].

In studies of human prion diseases with apes and monkeys, the use of one or two, or rarely three animals as recipients for a single inoculum has presented a significant problem

in evaluating the transmissibility of a particular inoculum from an individual patient. The transgenic mice contain a chimeric prion protein gene, e.g., Tg(MHu2M) mice, and hybrid mice e.g., Tg(HuPrP)^{Prnp} described here obviate many of the problems created by using nonhuman primates.

These results demonstrate the "universality" of the MHu2M transgene for transmission studies with other types of transgenic animals and other prion inocula. For example, it may be most efficient to use mice expressing MHu2MPrP transgenes coding for either a methionine or valine at codon 129, and by doing so, match the genotype of the Tg mouse (with respect to codon 129) with the genotype of the individual from which the inoculum is derived. Homozygosity at the codon 129 polymorphism has a profound influence on the incidence of sporadic CJD [Palmer et al., *Nature* 332:340-342 (1991)]. The MHu2MPrP transgene encodes a Met at codon 129 and the iatrogenic CJD case was homozygous for Met [Collinge et al., *Lancet* 337:1441-1442 (1991)].

A human PrP gene is polymorphic at codon 129. More specifically, normal human PrP gene can be either homozygous Met/Met or Val/Val or heterozygous Met/Val at codon 129. The codon 129 polymorphism influences the susceptibility of humans to prion disease and specifically to iatrogenic and sporadic CJD. This polymorphic codon is contained in the central region of MHu2MPrP which is derived from human PrP. The DNA sequence used to generate Tg(MHu2M) mice encodes Met at codon 129. The transgenic mice expressing MHu2MPrP with valine at codon 129 can be produced using similar procedures.

To break the species barrier an artificial PrP gene was inserted into a host mammal (such as a mouse) renders that mammal susceptible to infection with prions which normally infect only a genetically diverse test mammal (e.g. a human, cow or sheep). The artificial PrP gene may include the natural PrP gene sequence of the host animal with one or more (preferably less than 40) codon sequences being replaced with other codon sequences such as corresponding codons of a genetically diverse mammal (e.g. a human, cow or sheep).

The species barrier is broken by inserting into a mammal (a mouse) the chimeric gene (MHu2M) which is shown being assembled schematically in Figure 1. In order to produce the chimeric gene, it is first necessary to obtain nucleotide sequences which encode human PrP. The human PrP genes are then subjected to the conventional PCR procedures in order to produce large numbers of copies of the gene or portions of the gene. The PCR product is then isolated, specific restriction sites are added and the copied product is

subjected to specific endonucleases in order to remove a middle section of the human PrP gene. Specifically, restriction sites are added such that when the PCR product is subjected to endonucleases such as Asp718 as well as BstEII, a section of the gene is cut out. The use of these two endonucleases will remove a center portion of the human PrP gene (codons 94-188) which portion encodes amino acid residues 94 through 188. Endonucleases are also used to remove a corresponding center portion of the mouse PrP gene. The removed center portion of the mouse gene is then discarded and the center portion obtained from the human PrP gene is fused into the mouse gene to produce a chimeric human/mouse gene. Details of how the specific MHu2M gene was produced are described in Example 1 and shown in Figure 1.

As shown with Figure 2, there is a high degree of homology between the removed center portion of the human PrP gene and the segment of the mouse PrP gene which is replaced. Specifically, the segments differ at nine codons. Thus, when the genetic material is expressed, the resulting chimeric MHu2M protein will differ from MoPrP at 9 residues. These residues and their positions are shown in Figure 2. After the chimeric gene is produced, it can be microinjected into a mouse egg using known technology as described within Scott et al., *Cell* 59:847-857 (1989) and Scott et al., *Protein Sci.* 1:986-997 (1992) and see also WO91/19810 published December 22, 1991 as well as other publications relating to the production of transgenic mice cited therein and known to those skilled in the art. The injected mouse egg is then implanted into a mouse using known procedures. Multiple eggs can be implanted into a single mouse and known procedures can be used to determine whether the resulting offspring are transgenic mice which include the chimeric gene within their genome. Details of this procedure are described in Examples 6 and 7.

The "species barrier" is broken by producing a chimeric PrP gene wherein a middle portion of the mouse PrP gene is replaced with a corresponding middle portion of a human PrP gene, thereby leaving the C- and N-terminus of the mouse PrP gene intact. However, other segments of the mouse PrP gene can be replaced with other homologous segments of the human PrP gene and obtain a transgenic mouse which is subject to being readily infected with human prions.

PATHOGENIC MUTATIONS AND POLYMORPHISMS

There are a number of known pathogenic mutations in the human PrP gene. Further, there are known polymorphisms in the human, sheep and bovine PrP genes. The following is a list of such mutations and polymorphisms:

5

MUTATION TABLE

	Pathogenic human mutations	Human Polymorphisms	Sheep Polymorphisms	Bovine Polymorphisms
	2 octapeptide insert	Codon 129 Met/Val	Codon 171 Arg/Gln	5 or 6 octapeptides
10	4 octapeptide insert	Codon 219 Glu/Lys	Codon 136 Ala/Val	
	5 octapeptide insert			
	6 octapeptide insert			
	7 octapeptide insert			
	8 octapeptide insert			
15	9 octapeptide insert			
	Codon 102 Pro-Leu			
	Codon 105 Pro-Leu			
	Codon 117 Ala-Val			
	Codon 145 Stop			
20	Codon 178 Asp-Asn			
	Codon 180 Val-Ile			
	Codon 198 Phe-Ser			
	Codon 200 Glu-Lys			
	Codon 210 Val-Ile			
25	Codon 217 Asn-Arg			
	Codon 232 Met-Ala			

In order to provide further meaning to the above chart demonstrating the mutations and polymorphisms, one can refer to the published sequences of PrP genes. For example, a chicken, bovine, sheep, rat and mouse PrP gene are disclosed and published within Gabriel et al., *Proc. Natl. Acad. Sci. USA* 89:9097-9101 (1992). The sequence for the Syrian hamster is published in Basler et al., *Cell* 46:417-428 (1986). The PrP gene of sheep is published by Goldmann et al., *Proc. Natl. Acad. Sci. USA* 87:2476-2480 (1990). The PrP gene sequence for bovine is published in Goldmann et al., *J. Gen. Virol.* 72:201-204 (1991). The sequence for chicken PrP gene is published in Harris et al., *Proc. Natl. Acad. Sci. USA* 88:7664-7668 (1991). The PrP gene sequence for mink is published in Kretzschmar et al., *J.*

5 mice which develops symptoms of prion disease at the same time are sacrificed and their brains extracted and homogenized such will create a relatively standard prion preparation. This preparation can then be used to inoculate a new group of transgenic animals. This process can be repeated a number of times e.g., 1 to 10 times or until such point as all of the transgenic mice are developing symptoms of prion disease at approximately the same point in time after inoculation with the standardized preparation. Further details of how to produce a standardized preparation are provided below and in Example 8.

10 In that the preparation is to be a "standard" it is preferably obtained from a battery (e.g., 100 or 1,000 or more animals) of substantial identical animals. For example, 100 mice all containing a very high copy number of human PrP genes (all polymorphisms and mutations) would spontaneously develop disease and the brain tissue from each could be combined to make a useful standardized prion preparation.

15 Standardized prion preparations can be produced using any of the modified host mammals of the present invention. For example, standardized prion preparations could be produced using mice, rats, hamsters, or guinea pigs which are genetically modified per the present invention so that they are susceptible to infection with prions which prions would generally only infect genetically diverse species such as a human, cow, sheep or horse and which modified host mammals will develop clinical signs of CNS dysfunction within a period of time of 350 days or less after inoculation with prions. The most preferred host mammal is a mouse in part because they are inexpensive to use and because a greater amount of experience has been obtained with respect to production of transgenic mice than with respect to the production of other types of host animals.

20 Once an appropriate type of host is chosen, such as a mouse, the next step is to choose the appropriate type of genetic manipulation to be utilized to produce a standardized prion formulation. For example, the mice may be mice which are genetically modified by the insertion of a chimeric gene of the invention. Within this group the mice might be modified by including high copy numbers of the chimeric gene and/or by the inclusion of multiple promoters in order to increase the level of expression of the chimeric gene.

25 Alternatively, hybrid mice of the invention could be used wherein mice which have the endogenous PrP gene ablated are crossed with mice which have a human PrP gene inserted into their genome. There are, of course, various subcategories of such hybrid mice. For example, the human PrP gene may be inserted in a high copy number and/or used with

5 Gen. Virol. 73:2757-2761 (1992). The human PrP gene sequence is published in Kretschmar et al., DNA 5:315-324 (1986). The PrP gene sequence for mouse is published in Loch et al., Proc. Natl. Acad. Sci. USA 83:6372-6376 (1986). The PrP gene sequence for sheep is published in Westaway et al., Genes Dev. 8:959-969 (1994). These publications are all incorporated herein by reference to disclose and describe the PrP gene and PrP amino acid sequences.

DIFFERENCES IN THE CONVERSION OF MHu2MPp^{Sc} AND HuPrP^{Sc} INTO THE SCRAPIE ISOFORM IN MICE

10 The fundamental event in prion propagation is the conversion of PrP^C, which contains ~43% α -helix and is devoid of β -sheet, into PrP^{Sc}, which has ~44% β -sheet [Pan et al., Proc. Natl. Acad. Sci. USA 90:10962-10966 (1993)]. From the results of Tg(SHuPrP) mouse studies, this process is thought to involve the formation of a complex between PrP^{Sc} and the homotypic substrate PrP^C [Prusiner et al., Cell 63:673-686 (1990)]. Attempts to mix PrP^{Sc} with PrP^C have failed to produce nascent PrP^{Sc} [Raether et al., J. Virol. 66:6155-6163 (1992)], showing that PPMF is catalyzing the conformational changes that feature in the formation of PrP^{Sc}. The difference in susceptibility of Tg(MHu2M) and Tg(HuPrP) mice to Hu prions in mice is due to the fact that mouse PPMF catalyzing the refolding of PrP^C into PrP^{Sc} can recognize MHu2MPp much more readily than HuPrP.

STANDARDIZED PRION PREPARATION

20 Standardized prion preparations are produced for use in assays so as to improve the reliability of the assay. Although the preparation can be obtained from any animal it is preferably obtained from a host animal which has brain material containing prions of a test animal. For example, a Tg mouse containing a human prion protein gene can produce human prions and the brain of such a mouse can be used to create a standardized human prion preparation. The preparation can be further standardized by repeating the above process. More specifically, per the above process some prion containing material may be used to inoculate the transgenic mice. The source of that prion containing material may itself be unpredictable and result in infecting transgenic mice in different ways. Thus, if the transgenic mice are infected with a nonstandard material some may develop the symptoms of prion disease at different rates and some may not develop symptoms at all. If a group of

multiple promoters to enhance expression. In yet another alternative the mice could be produced by inserting multiple different PrP genes into the genome so as to create mice which are susceptible to infection with a variety of different prions, i.e., which generally infect two or more types of test animals. For example, a mouse could be treated which included a chimeric gene including part of the sequence of a human, a separate chimeric gene which included part of the sequence of a cow and still another chimeric gene which included part of the sequence of a sheep. If all three different types of chimeric genes were inserted into the genome of the mouse the mouse would be susceptible to infection with prions which generally only infect a human, cow and sheep.

After choosing the appropriate mammal (e.g., a mouse) and the appropriate mode of genetic modification (e.g., inserting a chimeric PrP gene) the next step is to produce a large number of such mammals which are substantially identical in terms of genetic material related to prions. More specifically, each of the mice produced will include an identical chimeric gene present in the genome in substantially the same copy number. The mice should be sufficiently identical genetically in terms of genetic material related to prions that 95% or more of the mice will develop clinical signs of CNS dysfunction within 350 days or less after inoculation and all of the mice will develop such CNS dysfunction at approximately the same time e.g., within ± 30 days of each other.

Once a large group e.g., 50 or more, more preferably 100 or more, still more preferably 500 or more of such mice are produced. The next step is to inoculate the mice with prions which generally only infect a genetically diverse mammal e.g., prions from a human, sheep, cow or horse. The amounts given to different groups of mammals could be varied. After inoculating the mammals with the prions the mammals are observed until the mammals exhibit symptoms of prion infection e.g., clinical signs of CNS dysfunction. After exhibiting the symptoms of prion infection the brain or at least a portion of the brain tissue of each of the mammals is extracted. The extracted brain tissue is homogenized which provides the standardized prion preparation.

As an alternative to inoculating the group of transgenic mice with prions from a genetically diverse animal it is possible to produce mice which spontaneously develop prion related diseases. This can be done, for example, by including extremely high copy numbers of a human PrP gene into a mouse genome. When the copy number is raised to, for example, 100 or more copies, the mouse will spontaneously develop clinical signs of CNS dysfunction

-41-

and have, within its brain tissue, prions which are capable of infecting humans. The brains of these animals or portions of the brain tissue of these animals can be extracted and homogenized to produce a standardized prion preparation.

The standardized prion preparations can be used directly or can be diluted and tined in a manner so as to provide for a variety of different positive controls. More specifically, various known amounts of such standardized preparation can be used to inoculate a first set of transgenic control mice. A second set of substantially identical mice are inoculated with a material to be tested i.e., a material which may contain prions. A third group of substantially identical mice are not injected with any material. The three groups are then observed. The third group, should, of course not become ill in that the mice are not injected with any material. If such mice do become ill the assay is not accurate probably due to the result of producing mice which spontaneously develop disease. If the first group, injected with a standardized preparation, do not become ill the assay is also inaccurate probably because the mice have not been correctly created so as to become ill when inoculated with prions which become ill and the third group does not become ill the test material does not contain prions and if the second group does become ill the test material does contain prions.

By using standardized prion preparations it is possible to create extremely dilute compositions containing the prions. For example, a composition containing one part per million or less or even one part per billion or less can be created. Such a composition can be used to test the sensitivity of the transgenic mice of the invention in detecting the presence of prions in the sample.

Prion preparations are desirable in that they will include a constant amount of prions and are extracted from an isogenic background. Accordingly, contaminants in the preparations will be constant and controllable. Standardized prion preparations will be useful in the carrying out of bioassays in order to determine the presence, if any, of prions in various pharmaceuticals, whole blood, blood fractions, foods, cosmetics, organs and in particular any material which is derived from an animal (living or dead) such as organs, blood and products thereof derived from living or dead humans. Thus, standardized prion preparations will be valuable in validating purification protocols where preparations are spiked and reductions in titer measured for a particular process.

-42-

MEASURING LEVELS OF PRIONS

PPMF can be administered to transgenic mice to determine the concentration of prions (which generally only infect a genetically diverse animal) within a given sample. The transgenic mice make it possible to test for the positive presence of prions within a sample.

The mice are capable of detecting the presence of prions in a concentration as low as 1 ppn or even 1 ppb or less and administering PPMF reduces the time needed for the mice to show signs of disease. The procedure for doing such will be apparent to those skilled in the art upon a review of the present disclosure in combination with an article entitled "Measurement of The Scrapie Agent Using An Incubation Time Interval Assay," published by Prusiner, et al, *Annals of Neurology* 11:353-358 (1982) which is incorporated herein by reference to disclose such a method of measurement. See also U.S. Patent No. 5,565,186, issued October 15, 1996. In general, the method is carried out by determining the titer of the prions by carrying out measurements of time intervals from inoculation to onset of symptoms and from inoculation to death. The intervals are inversely proportioned to the size of the dose injected intracerebrally. The logarithms of the time intervals minus a time factor are linear functions of the logarithms of the inoculum size.

EVIDENCE OF DISEASE

PrP^{Sc} has been found in the brains of affected Tg(MHu2M) mice after inoculation with Hu(CJD) or Mo(RML) prions. Brain homogenates of Tg(MHu2M) were either left undigested or digested with proteinase K (BMB) at a final concentration of 20 µg/ml for 1 hour at 37°C (even numbered lanes). Samples were resolved by SDS PAGE and then analyzed by Western blot.

The distribution of PrP^{Sc} and PrP^{Sc} in clinically sick Tg(MHu2M) mice inoculated with Mo(RML) and Hu(CJD) prions were detected by the histoblot method. The histoblots included those of coronal sections through the region of the hippocampus and thalamus. Differences are observed between: (A) PrP^{Sc} in Mo(RML) infected mouse; (B) PrP^{Sc} in sporadic CJD RG-infected mouse; (C) PrP^{Sc} in Mo(RML) infected mouse; (D) PrP^{Sc} in sporadic CJD RG-infected mouse; (E) PrP^{Sc} in sporadic CJD EC-infected mouse; and (F) PrP^{Sc} in iatrogenic CJD (#364)-infected mouse.

STRUCTURAL ANALYSIS OF PrP^{Sc}

To investigate the basis of the PrP structural transitions, recombinant PrP protein, termed rPrP, was prepared using the SHaPrP sequence corresponding to residues 90-231 of PrP 27-30. The rPrP was uniformly labeled with ¹⁵N or with both ¹³C and ¹⁵N; it was refolded into a conformer that resembles PrP^{Sc} based on optical spectroscopic and immunochemical measurements. The pH of the rPrP solution was found to be critical. N-terminal epitopes (residues 90-112) in rPrP were observed to be partially buried at pH 5.2 - 8.0 by ELISA using N-terminal FfAbs. These epitopes became completely exposed at pH 4.8 or lower. Whether this conformational transition, detected by FfAbs, modified the NMR structure of MoPrP (121-231) determined at pH 4.5 (3) remains to be established. When rPrP (0.9 mM) was poised in the middle of this structural transition at pH 5.0 in 20 mM acetate and 0.005% Na azide, its α-helical state was stable for at least 15 days at temperatures from 4-30°C as judged by circular dichroism (CD); however, even one day at 35°C led to a significant loss of α-helix. This apparent acquisition of β-sheet was concentration-dependent: at 35°C, increasing the concentration from 0.03 mM incrementally to 0.75 mM revealed that increasing the protein concentration steadily increased the rate of the conversion.

Multidimensional heteronuclear NMR studies were performed with rPrP. Signal line widths and spectral dispersion indicated that most of the protein is well-structured with the concentration (ca. 1 mM) and solution conditions employed: 20 mM sodium acetate, pH 5.2, 30°C. Analytical sedimentation confirmed that the protein was essentially monomeric; samples reached equilibrium after 15 h and did not change between 15 and 40 h at 25°C. The three sets of data fit well with a monomeric molecular weight of 16, 242 kDa with small residuals. Assuming a monomer-dimer equilibrium, the association equilibrium constant was calculated to be $5.4 \times 10^6 \text{ M}^{-1}$. These findings indicate that refolded rPrP is predominantly monomeric under the conditions of the NMR experiments.

Resonance assignments were made using samples of uniformly ¹⁵N- and ¹³C- ¹⁵N- labeled rPrP, primarily via scalar-coupling-based three-dimensional NMR spectra, including total correlated spectroscopy (TOCSY). There are essentially three parts to the protein really reflected in simple spectral features: residues 90-112 are characterized by relatively narrow ¹⁵N HSQC spectral signals and many NOE cross-peaks; residues 113-126 have

relatively narrow ^1H HSQC spectral signals and many NOE cross-peaks; and most of the remaining residues exhibit *ca.* 6z broader HSQC signals and numerous NOE cross-peaks.

The α -proton and α -carbon chemical shift indices, as well as the proton NOE connectivities evidence in nuclear Overhauser effect spectra (NOESY) consistently indicate that rPrP contains three α -helical regions. The locations of these correspond largely, but not entirely, to those found for MoPrP (121-231) under similar solution conditions (0.8 mM protein, pH 4.5, no buffer, 20°C) and to two of the four helices predicted for the entire sequence.

From the NOE cross-peaks, 2401 experimental distance restraints were used to generate low-resolution structures via the program DIANA, followed by minimization with AMBER 4.1. A best-fit superposition of backbone atoms for residues 113-228 of rPrP is shown in Figure 5. The helices and β -sheet are fairly well-defined, and loop regions can also be defined, although with less precision. To distinguish the α -helices found in rPrP by NMR from those predicted by molecular modeling, these were provisionally designated helices A, B and C. Helix A spans residues 144-157 with the last turn being quite distorted, corresponding to helix 144-154 found for MoPrP (121-231). Helix B spans residues 172-193, with the first turn being irregular at the present state of structure refinement. This is about two turns longer than the 179-193 helix found for MoPrP (121-231) which agrees well with predicted helix H3 (179-191). Helix C extends from residue 200 to 227 with the 225-227 turn being irregular, which is about three turns longer than the helix corresponding to residues 200-217 in MoPrP (121-231). It is notable that predicted helix H4 (residues 202-218) corresponds well with that found in MoPrP (121-231). Two four-residue β -strands (128-131 and 161-164) were identified in the MoPrP (121-231) structure. A similar antiparallel β -sheet was found, with S2 spanning residues 161-163 and S1 spanning 129-131 possessing β -sheet characteristics, but the two strands do not manifest standard β -sheet geometry. In fact, a β -bridge occurs only between Leu¹⁴⁶ and Tyr¹⁴², and there are extensive cross-strand connectivities of residues in segment 129-134 with proximate residues on the antiparallel segment 159-165.

The loop between S2 and helix B (i.e., residues 165-71) yields resonances clearly exhibiting long-range as well as medium-range restraints in contrast to the absence of resonances for the backbone atoms of residues 167-176 in the shorter MoPrP (121-231). These results indicate that the loop is reasonably ordered, whereas it was concluded that this

region is disordered in MoPrP (121-231). Figures 6A and 6B show only one example: ^1H - ^1H cross-peaks between the unresolved methyl protons of Val¹⁶⁶ in that loop with Ser²²².

However, the methyl protons of Val¹⁶⁶ also exhibit 15 long-range cross-peaks with protons in the same loop, e.g., Tyr¹⁴⁹, and in the extension of helix C, e.g., Tyr²³¹ and Tyr²²¹, which are on the same side of the helix. Connectivities of Val¹⁶⁶ to residues two turns apart on helix C suggest that the loop exits in multiple discrete conformations. In this context, it is noteworthy that the loop is not under-defined, indeed, the multiple observed connectivities over-define the loop. Apparently, the interaction of the 165-171 loop with the helix C extension is important in stabilizing the structure. Figure 5 illustrates the relationship of the loop to helix C.

The α -proton and α -carbon chemical shifts for residues 90-127 are consistent with the region having substantial α -helical content, but the extent of the chemical shifts relative to that of random coil values was generally not enough to indicate the α -helix formation via tripartite chemical shift indices. There are also insufficient NOE connectivities to conclude α -helix is formed. The few medium-range connectivities in the segment 90-112 demonstrate sparse elements of structure. For example, for residues 95-100, results have so far identified 9 nonsequential NOE connectivities which imply that significant structure exists at least transiently. The small number of long-range connectivities for the N-terminal segment 90-112 imply that it is largely disordered.

Results obtained have identified 36 long-range NOE cross-peaks involving sidechain resonances for the hydrophobic residues in the segment 113-125. An uncommon combination of glycines and hydrophobic residues leads to an unusual, and dynamic, structural feature. Most of the NOE connectivities indicate that these residues form a hydrophobic globule with substantial backbone reversals permitted by the many glycines;

indeed Val¹²¹, Val¹²², and Leu¹²³ each exhibit 10-13 long-range connectivities. As seen in Figure 5, the backbone for this globule is not well-defined in spite of any connectivities. This may reflect the true dynamic nature of such a hydrophobic globule, as the ^1H HSQC spectral line widths for these residues were also about 6 Hz smaller than for the core of the protein (*ca.* 18 vs. 24-25 Hz). Apparently, the combination of glycines with hydrophobic residues permits many alternative conformations with comparable free energies.

Some long-range connectivities place the hydrophobic globule adjacent to the β -sheet in contact with the S1 strand (Figure 7). The weak, broadened ^1H HSQC spectral signals for

S1 residues Met¹²⁹ (32 Hz), Leu¹³⁰ (32 Hz) and Gly¹³¹ (44 Hz) as well as S2 residues Tyr¹⁶² (33 Hz) and Arg¹⁶⁴ (33 Hz) may well reflect conformational exchange effects in the interacting hydrophobic globule pervading the adjacent irregular β -sheet. Taken together, these results suggest that the hydrophobic globule and adjacent β -strands constitute a domain with marginally stable polymorphic structure.

The apparent conformational heterogeneity of the N-terminal region of rPrP may reflect the process by which PrP^C is converted into PrP^{Sc}. Transgenic studies also show that PrP^{Sc} formation requires the substrate PrP^C to bind the product PrP^{Sc} at an intermediate stage of the conversion process. PrP^C is thought to be in equilibrium with a metastable intermediate, designated PrP^I, which binds to PrP^{Sc} in the conversion process. In fact, destabilization of PrP^C has been shown to be necessary for it to bind to PrP^{Sc} *in vitro*. Further evidence for the conformational plasticity of PrP comes from unfolding studies of rPrP using guanidinium chloride (GdnHCl). The free energy difference ΔG_3 of 6.5 ± 1.2 kcal/mole between an intermediate state and the unfolded state was found to be comparable to literature values ($5-15$ kcal/mole) for protein unfolding. However, the completely refolded rPrP, as used for the present NMR studies, is only marginally more stable ($\Delta G_1 = 1.9 \pm 0.4$ kcal/mole) than the folding intermediate. This is consistent with the extensive conformational flexibility evident in the current NMR studies for part of the protein. The results also show another α -helical form of rPrP, formed by refolding at pH2, which exhibits NMR features characteristic of an acid-denatured molten globule but one that largely converts to the isoform studied here upon sitting for a few days at pD5.4 (corresponding to a pH meter reading of 5.0 measured in H₂O).

The NMR results for rPrP, compared to the structure reported for MoPrP, support the notion that the core of the PrP^{Sc} structure is formed by parts of helices B and C, corresponding largely to the predicted H3 and H4 regions, and stabilized by the disulfide which is essential for α -helical folding. As seen in Figures 8 and 9, helices B and C essentially form one side of the protein structure. This core is further stabilized by helix A, which lies across helix C with sidechains between the two helices interacting (Figure 8). Strand S2 also lies on this side of the protein and interacts predominantly with helices B and C as well as S1. With or without S2 and S1, we presume this relatively stable folding core is associated with the second unfolding transition. Attempts to prepare MoPrP (108-231) resulted in proteolytic cleavage producing MoPrP (121-131), indicative of a stable core

beginning with residue 121. Conclusions about the "stable core" of PrP, however, must be considered within the context of a mutagenesis studies of a PrP with residues 23-89 and 141-176 deleted, i.e. removing helix A and strand S2, the resulting 106-residue PrP could be converted to a protease-resistant form.

Presence of the additional 31 N-terminal residues of rPrP, relative to MoPrP (121-231), induces substantial changes in the structure of PrP including alterations in the C-terminus. Helix C is extended by at least nine residues, helix B is up to seven residues longer, and the loop compromising residues 165-173 is sufficiently ordered that many long-range restraints can be observed. The hydrophobic globule (residues 113-125) predominantly interacts with S1 in the β -sheet (Figure 8). This may serve to stabilize the observed extension of helix B from 179 in MoPrP (121-231) to 172 in rPrP. Stability may also be conferred by hydrophobic interactions of Tyr¹²³ with Tyr¹⁶³ in the β -sheet which, in turn, interacts with Val¹¹⁹. The relative stability of the 165-171 loop and the three additional helical turns in helix C are presumably connected to stabilization of the other structural elements.

Within the hydrophobic globule is a palindromic sequence, A¹¹³GAAAGA, that is conserved in all species examined to date, as are the surrounding residues. In humans the A117V mutation causes GSS, while an artificial set of mutations consisting of A113V, A115V, and A118V in Tg mice causes spontaneous neurodegeneration, promotes β -sheet formation in recombinant PrP, and generates prion infectivity *de novo*.

Strains of prions exhibit different incubation times before symptoms of disease appear and different patterns of PrP^{Sc} accumulation. Recent work indicates that the properties of prion strains might be manifestations of different conformers of PrP^{Sc}. See Telling, et al, Science, Dec. 1996. Studies on the transmission of human prions to transgenic mice also show that a species-specific PP2MF, functions as a molecular chaperon in PrP^{Sc} formation. These results show that PP2MF forms a transient complex with the metastable intermediate PrP, diminishing the activation energy barrier between PrP^C and PrP^{Sc} and facilitating formation of PrP^{Sc}. An analysis comparing the three-turn extension of helix C to residue 227 in rPrP with the helix termination at Gly²¹⁷ in MoPrP(121-231) is consistent with this concept. SHa residues Tyr²¹⁵ and Glu²¹⁹ are at the site of PP2MF binding and the glycosylation sites, Asn¹¹⁸, and Asn¹¹⁹, are not very near this binding site (Figure 9). In this context, the stability and length of helix C become an important issue. Interestingly,

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

EXAMPLE 1

BINDING SITE IN PrP^C for PPMF

An examination of the amino acids which distinguish Hu PrP from Mo PrP shows only seven residues at the C-terminus (168-231) that are different. Figure 3 is a comparison of sequences showing the differences. Four of these residues are close to the glycoprophosphatidylinositol (GPI) anchor attached to Ser231 while the remaining three residues were within or near the C-terminus of a postulated α -helix which has been conformed by NMR structural studies. To identify the critical binding site within PrP^C for PPMF the seven residues were divided into two groups: those at the C-terminal end of the last α -helix (HuA) and those at the extreme C-terminus (HuB) (Figure 3). The Mo residues were replaced with Hu counterparts in positions that were critical for binding of PrP^C to Mo PPMF to determine the effect of such on inhibiting the formation of recombinant PrP^{Sc}. Recombinant PrP^{Sc} was distinguished from endogenous wild-type (wt) Mo PrP^{Sc} by using the SHa/Mo chimeric PrP designated MHMHuA that contains a binding site for the anti-SHa PrP 3F4 monoclonal antibody (mAb).

Three chimeric constructs, denoted as MHMHuA (Mo residues 214, 218 and 219 were replaced with the corresponding human residue), MHMHuB (Mo residues 226 through 230 replaced with Hu), and MHMHu(A+B) (combined replacements), were transiently transfected into scrapie infected Mo neuroblastoma (ScN2a) cells (Figure 3). Neither MHMHu(A+B) nor MHMHuA was converted into PrP^{Sc} as judged by the acquisition of protease resistance. By contrast, MHMHuB was converted into PrP^{Sc} as efficiently as the

Residues where point mutations lead to human diseases are highlighted in Figure 9.

A point mutation in the PrP gene encoding Asp¹⁷⁸ leading to Asn¹⁷⁸ causes fatal familial insomnia if residue 129 is Met; the double mutation with Met¹²⁹ mutated to Val as well results in a subtype of CJD instead. Residue 178, which is in the extension of helix B seen in rPrP but not in MoPrP (121-231), and residue 129 are located opposite one another with strand S2 partially intervening (Figure 9). If the mutation D178N destabilizes the structure, part of helix B could unravel. This part of helix B is near Arg¹⁶⁴ in S2, which in turn is adjacent to Met¹⁷⁹. Depending upon the identity of residue 129, the structure (or, in light of its stability, structural ensemble) resulting from this destabilization may differ.

Our structural studies of rPrP underscore the conformational plasticity evident in the N-terminal region and defines important structural features not evident in a smaller C-terminal fragment. Prior studies, including most recently with F1abs, indicate that the region corresponding to the N-terminal 30-40 residues of rPrP change conformation during formation of PrP^{Sc}.

control MHM2. These results indicate that Mo PPMF did not bind to MHMHuA(+B) or to MHMHuA but did bind to MHMHuB and MHM2, both of which were converted into Pp^{sc}.

EXAMPLE 2

Example 1 identifies the HuA region which prevents conversion of modified Pp^{sc} into Pp^{sc}. Additional constructs were then produced with Mo residue 214 replaced by its Hu counterpart (MHMHuA1), 218 and 219 replaced by the Hu homologues (MHMHuA2), Mo 218 replaced by its Hu counterpart (MHMHuA3), Mo 219 replaced by the Hu one (MHMHuA4), and Mo 214 and 218 replaced by the Hu homologues (MHMHuA5) (Figure 3). Neither MHMHuA2, MHMHuA3 nor MHMHuA5 were converted into Pp^{sc}, while MHMHuA1 was converted but at a reduced level compared to MHMHuA4 and MHMHuB. It should be noted that longer film exposure revealed what appeared to be a trace amount of Pp^{sc} formation in MHMHuA2 and MHMHuA3 but not in MHMHuA5.

The experiments described above show that substitution of the Hu encoded Glu residue for Gln in Mo Pp^{sc} at position 218 prevented its conversion into Pp^{sc}. In addition, substitution of the Hu encoded Ile residue for Val in Mo Pp^{sc} at position 214 ameliorated this inhibition. These data argue that the region of Pp^{sc} around the segment bounded by residues 214 and 218 participates in the formation of Pp^{sc}. There is no known metal, protein, or DNA binding motif in this region. Experiments with phosphatidylinositolphospholipase C (PIPLC) revealed no topological changes in these chimeric constructs.

EXAMPLE 3

ARTIFICIAL MUTATIONS AT Mo RESIDUES 214 and 218

A Hu polymorphism at residue 219 (corresponding to Mo 218) has been reported in the Japanese population where about 12% of the people have the Lys allele instead of the Glu/Gln allele. Experiments were carried out to investigate the influence of the polymorphism by introducing Lys at Mo residue 218 into MHM2 Pp^{sc} (MHMK 218, Figure 3). The results obtained show that Lys at Mo 218 completely abolished the Pp^{sc} formation (Table 1). This result indicates that the Lys allele at Mo 218 has an aberrant affinity to Mo PPMF and subsequently exhibits no Pp^{sc} formation. To further investigate the effect of amino acid substitutions at the residue 218, four artificial mutations were introduced; Ile

(MHM1218), Ala (MHMA218), Trp (MHMW218), Pro (MHMP218) (Figure 3). MHM1218 and MHMW218 converted into Pp^{sc} but much less efficiently, whereas neither MHMA218 nor MHMP218 showed conversion to Pp^{sc}.

Mutations at residue 214 hindered the formation of recombinant Pp^{sc} but in a different manner. In addition to MEMHuA1 (Ile at 214), MHMA214 (Ala at 214) was converted, albeit less efficiently, into MHMA214 Pp^{sc} however MHMK214 (Lys at 214), MHME214 (Glu at 214) MHMW214 (Trp at 214), nor MHMP214 (Pro at 214) were not converted (Table 1). These experiments show that amino acid 214 is on the periphery of the binding interface between Pp^{sc} and PPMF while residue 218 occupies a more central region of that binding interface.

EXAMPLE 4

BINDING CHARACTERISTICS OF PPMF TO Pp^{sc} IN Pp^{sc} FORMATION

In order to determine how residue 218 provides species specific binding to PPMF, mutants of Pp^{sc} were serially co-transfected into ScN2a cells to analyze any interactions between the different Pp^{sc} and Mo Pp^{sc} under conditions where the concentration of PPMF can be limiting. In this regime, inhibition is observed between Mo and SHa Pp in ScN2a cells in a manner similar to that observed in Tg mice showing significantly reduced Mo Pp^{sc} following Mo prion inoculation in the presence of heterologous expression of SHa Pp^{sc}.

Since neither SHa nor MH3 Pp^{sc} (Figure 3) could be efficiently converted into Pp^{sc}, MHM2 Pp^{sc} formation was dramatically reduced when co-transfected. MH2M Pp^{sc} (Figure 3) was less efficiently converted into Pp^{sc} with an intermediate amount of inhibition on MHM2Pp^{sc} formation when co-transfected. On the other hand, co-expression of Mo Pp^{sc} (Figure 3), which was efficiently converted into Pp^{sc} (JF4 does not recognize Mo Pp^{sc}), did not affect MHM2 Pp^{sc} formation when co-transfected.

Neither MH3HuA, MH2HuA Pp, nor M3HuA Pp exhibited any inhibition on MHM2 Pp^{sc} formation when co-transfected even if they did not convert into Pp^{sc} (Table 1) in marked contrast to SHa or MH3 Pp^{sc}. When the HuA sequence was introduced into MH3 Pp^{sc} (MH3HuA), the inhibition with SHa derivatives was abolished. This indicates that the HuA region primarily governs the inhibition in the Pp^{sc} formation over the Pp^{sc}-Pp^{sc} interaction determined by the central region.

Mutant PrP^{Sc} failed to efficiently convert into the PrP^{Sc} isoform, indicating that mutations at Mo residue 218 might produce a tightly associating complex between PP2F and PrP^{Sc} thereby eliminating subsequent PrP^{Sc} formation. To confirm such constructs carrying each mutation at residue 218 were co-transfected with MHM2 PrP^{Sc}. The higher the binding affinity between PP2F and the mutant protein the more unbound PP2F is depleted resulting in suppression of the conversion of co-expressed PrP^{Sc} into PrP^{Sc}. In addition, MHMHuA derivatives (Glu at 218), MHM1218 as well as MHM218 demonstrated no significant inhibition on the formation of MHM2 PrP^{Sc} when co-transfected. On the other hand, MHMA218 as well as MHMW218 exhibited remarkable inhibition (Table 1). These results indicate that MHMK218, MHMA218 and MHMW218 bind to PP2F even tighter than wt-MH2 (Glu at 218) does.

At residue 214, MHM1214, MHME214 and MHMW214 exhibited competitive inhibition on the conversion of MHM2 PrP^{Sc} into PrP^{Sc} when co-transfected, whereas MHMK214, MHMA214 and MHMP214 did not (Table 1). At the Mo residue 214, MHM1214, MHIE214, and MHMW214 have tighter binding to PP2F than MHM2-PrP^{Sc} does.

EXAMPLE 5

MUTATIONS AT Mo RESIDUE 216

A Swedish family with GSS has been reported with a mutation of Glu 217 Arg (216 in Mo). Brain samples in this GSS family showed no infectivity or protease resistant PrP^{Sc} on Western blots. To determine binding affinity to PP2F, Arg was introduced at Mo residue 216 in MHM2 PrP carrying either the Mo (MHMR216) or Hu sequence at residue 218 (MHMHuR216). Neither MHMR216 nor MHMHuR216 produced any PrP^{Sc} on Western blots (Table 1). When co-transfected with MHM2 PrP^{Sc}, however, MHMR216 (Glu at 218) inhibited formation of the MHM2 PrP^{Sc}, while MHMHuR216 (Glu at 218) did not (Table 1). This limited set of mutations at Mo residue 216 shows that binding of PP2F to PrP^{Sc} is independent of residue 216. This is consistent with the quite distinct orientations of side chains of residue 216 and 218 on the C-terminal α -helix of PrP^{Sc}.

30

MATERIALS AND METHODS

CULTURED CELLS

N2a cells were obtained from American Tissue Culture Collection (Rockville, MD). ScN2a cells are the persistently infected clones as described [Butler, et al., *J. Virol.* 62:1558-1564 (1988)]. All the cells were grown and maintained at 37°C in MEM supplemented with 10% FBS. In some cases, cells were treated with PIPLC. Cells grown on 60 mm dishes were rinsed with ice-cold PBS, then incubated with 0.5 U PIPLC in 1 ml of OptiMEM at 37°C for 4 h with swirling.

ANTIBODIES

α -PrP 3F4 is a mAb raised against SHa PrP27-30 [Kascsak, et al., *J. Virol.* 61:3688-3693 (1987)]. To distinguish the signals of MHM2-constructs from endogenous Mo PrP^{Sc}, 3F4 mAb was used since this antibody exclusively recognizes SHa, MHM2 PrP, MH2M, and MH3 PrP derivatives at Met¹⁰⁹-Met¹¹¹ [Rogers, *Glycobiology* 1:101-109 (1990)]. R073 is an antiserum raised in a rabbit against SDS-PAGE purified SHa PrP 27-30 reacts with SHa MH3, MH2M, M3H, Mo, and MHM2 PrP derivatives [Serban, et al., *Neurology* 40:110-117 (1990)].

RECOMBINANT GENE CONSTRUCTION

MH2M PrP was constructed as described [Scott, et al., *Protein Sci.* 1:986-997 (1992)]. MHM2 PrP in pSPOX was digested with BstEII and HindIII, then the 1.4 kbp fragment was ligated into BstEII-HindIII double digested Hu PrP in pSPOX, yielding MHM2Hu(A+B). MHMHu(A+B) in pSP72 was digested with XhoI and SmaI, then either the 2.4 kbp or the 880 bp fragment was ligated to MHM2 in pSP72, yielding MHMHuA or MHMHuB, which were then introduced into the expression plasmid pSPOX.

MH2M or Mo PrP in pSPOX was digested with BstEII and HindIII, then 1.4 kbp fragment was ligated into BstEII-HindIII double digested MHMHuA PrP in pSPOX, yielding MH2HuA and M3HuA. BstEII and XhoI restriction sites were introduced in the MHM2 by using the mismatched (underlined) BstH (5'ATC AAG CAG CAC ACG GTC ACC ACC ACC AAG GGG GAG AAC TTC ACC GAG ACC GAT ATC AAG ATA ATG) resulting in the mutations Val₂₀₈Ile and Met₂₀₉Ile, coupled with Xho (5' CAC TAT AGA ACT CGA GCA GCC TOC CT). After the PCR, amplified fragments were digested with restriction

WO 98/5132
PCT/US98/0104

their brains were taken for histopathologic studies (as per the procedures of Example 8) and confirmation of the diagnosis of scrapie.

EXAMPLE 11

IMMUNOBLOT ANALYSIS

For the determination of the relative levels of PrP expression in Tg mouse and human brains, protein concentrations were determined by bicinchoninic acid assay and immuno dots blot as previously described [Scott et al., *Cell* 73:979-988 (1993)]. Samples for Western blot analysis were prepared and western blots were performed as described previously [Towbin et al., *Proc. Natl. Acad. Sci. USA* 76:4350-4354 (1979)], except that an enhanced chemiluminescent (ECL) detection method (Amersham) was used. The blot was exposed to X-ray film for 5-60 seconds. α -PrP R073 rabbit antiserum was used at a final dilution of 1:5000 and 3F4 monoclonal antibody was also employed [Serban et al., *Neurology* 40:110-117 (1990)].

EXAMPLE 12

CREATING MAMMALS WHICH EXPRESS PPMF

Identifying the amino acid sequence(s) in MoPrP responsible for the binding of mouse PPMF and construct a modified MoPrP gene in which the sequence for this binding site is mutated. Such a benign MoPrP molecule will not interfere with human prion propagation in transgenic mice expressing HuPrP because PPMF is not sequestered by the mutant MoPrP. Following the above-described procedures in which the MoPrP gene is replaced with HuPrP or modified MoPrP genes, it is possible to create transgenic mice expressing HuPrP using these new genetic backgrounds either by genetic crosses or by direct microinjection of a vector capable of directing expression of HuPrP into fertilized embryos from these newly-created transgenic mouse lines.

EXAMPLE 13

MUTANT PrP CONVERSION INTO PrP^{Sc}

Three chimeric constructs, denoted as MHMHua (Mo residues 214, 218 and 219 replaced with Hu), MHMHuB (Mo residues 226 through 230 replaced with Hu), and MHMHu(A+B) (combined replacements), were transfected transiently into scrapie infected

-58-

WO 98/5132
PCT/US98/0104

Example 10 below, these mice were found to be susceptible to infection with human prions 100% of the time.

EXAMPLE 8

PREPARATION OF BRAIN HOMOGENATES

A 10% (w/v) homogenate of a sample of thawed human brain tissue was prepared in phosphate buffered saline lacking calcium and magnesium ions. The tissue was initially dissociated using a sterile disposable homogenizer, and this suspension was subjected to repeated extrusion through an 18 gauge syringe needle followed by a 22 gauge needle. Samples for inoculation into test animals were diluted 10-fold. Homogenates of clinically sick Tg and non-Tg mouse brains were prepared in the same way except for the omission of the initial dissociation step.

EXAMPLE 9

SOURCES OF PRION INOCULA

Human inocula were derived from frozen brain tissues of patients in which the clinical diagnosis of CJD or GSS had been confirmed by histopathological examination of brain tissues and, in most cases, by prion protein analysis. In some cases, the PrP gene was amplified by PCR of DNA isolated from patient blood and the PrP sequence determined by DNA sequence analysis. No HuPrP mutations were detected in cases of sporadic or iatrogenic CJD. The RML isolate was obtained from Swiss mice [Chandler, R.L., *Lancet* i:1378-1379 (1961)] from a closed colony at the Rocky Mountain Laboratory or in Swiss CD-1 mice obtained from Charles River Laboratories.

EXAMPLE 10

DETERMINATION OF SCRAPIE INCUBATION PERIODS

Transgenic mice as per Example 7 were inoculated intracerebrally with 30 μ l of brain extract using a 27 gauge needle inserted into the right parietal lobe. The preparation of inocula and criteria for diagnosis of scrapie in mice have been described [Carlson et al., *Cell* 46:503-511 (1986)]. Beginning 50 days after inoculation, the mice were examined for neurologic dysfunction every 3 days. When clinical signs of scrapie appeared, the mice were examined daily. When some animals whose death was obviously imminent were identified,

-57-

Mo neuroblastoma (ScN2a) cells. Neither MHMHu(A+B) nor MHMHuA was converted into Prp^{as} as judged by the acquisition of protease resistance. In contrast, MHMHuB was converted into Prp^{as} as efficiently as the control MHM2. We interpreted these results as indicating that Mo PPMF did not bind to MHMHu(A+B) or MHMHuA but did recognize MHMHuB and MHM2, both of which were converted into Prp^{as}. The mutant Prp molecules were all expressed at about the same level, and no inhibition of wt MoPrp^{as} formation could be detected.

Having identified the HuA region which prevents conversion of modified Prp^{as} into Prp^{as}, we produced additional constructs with Mo residues 214, 218 and 219 replaced by their Hu counterparts (Figure 10). To test the replacement of these residues either alone or in combination, we generated five constructs which were expressed then in ScN2a cells. Substitution of Hu residue 218 abolished Prp^{as} while substitution of Hu residue 219 was not inhibitory. Substitution of Hu residue 214 was partially inhibitory.

These results show that substitution of the Hu encoded Glu residue for Gln in Mo Prp^{as} at position 218 prevented its conversion into Prp^{as}. In addition, substitution of the Hu encoded Ile residue for Val in Mo Prp^{as} at position 214 was also inhibitory. These data argue that the region of Prp^{as} around the segment bounded by Mo residues 214 and 218 participates in the formation of Prp^{as}. Since Mo residues 214 and 218 lie one turn apart with the C-terminal α -helix of Prp, their side chains protrude from the same surface. These data argue that the region of Prp^{as} around the segment bounded by Mo residues 214 and 218 participates in the formation of Prp^{as}. There is no known metal, protein, or DNA binding motif in this region. Since the GPI anchor of Prp^{as} targets to caveolae where Prp^{as} formation seems to occur. An inquiry was made as to whether any of these amino acid substitutions prevented GPI anchor addition and transport of Prp^{as} to the cell surface. Studies of chimeric Prp^{as} released from the cell surface with phosphatidylinositolphospholipase C (PIPLC) digestion revealed no topological changes.

EXAMPLE 14

SPECIFICITY OF PrP MUTATIONS AT Mo RESIDUES 214 AND 218

Since substitution of the negatively charged Hu Glu residue for Mo Gln at position 218 inhibited conversion of this mutant Prp into Prp^{as}, experiments were carried out to determine if other amino acids would exhibit a similar effect. In humans, position 219

corresponding to mouse 218 is polymorphic: in Caucasians only Glu at this residue has been reported, whereas about 12% of Japanese people have the Lys allele. The substitution of Lys at Mo residue 218 abolished the Prp^{as} formation. To examine the specificity of amino acid substitutions at position 218, an examination was made of the effect of amino acid substitutions at the residue 218, by introducing 7 artificial mutations, Ile, Ala, Trp, Phe, Arg, or His (See Figure 3). The constructs expressing Ala, Pro, Phe, Arg, or His at position 218 were not converted into Prp^{as}, whereas, low amounts of Prp^{as} were made with the Ile or Trp at residue 218.

Substitution of Hu Ile at residue 214 diminished but did not completely abolish Prp^{as} formation. Similarly, modest Prp^{as} formation was observed with Ala. No Prp^{as} was observed when Lys, Glu, Trp or Pro were substituted at position 214. The mutant Prp molecules were all expressed at about the same level except when Pro was substituted at position 218, Arg at 216 or Pro 214. Substitutions of Pro in the C-terminal α -helix are expected to destabilize this secondary structure and may result in the increased lability of the protein. No inhibition of wt MoPrp^{as} formation by the mutant Prp molecules could be detected.

EXAMPLE 15

MUTANT PrPs AND CONVERSION OF WT PrP INTO PrP^{as}

Since only a minority of the ScN2a cells express the mutant PrPs in these transient transfection experiments, it was not possible to assess the effect of expressing mutant PrP conversion of wt MoPrP into Prp^{as}. To measure the influence of mutant PrP on the conversion of wt PrP into Prp^{as} cotransfection studies were performed. In these experiments, epitope tagged MHM2 PrP without (wt) or with a mutation were transfected together into ScN2a cells. Several studies have established that the transfecting DNAs are generally taken up together and coexpressed.

Substitution of Glu, Ile, Pro or Phe at residue 218 did not inhibit conversion of epitope tagged wt MHM2 Prp^{as} into Prp^{as}. In contrast, Lys, Ala, Trp, Arg or His at position 218 inhibited wt Prp^{as} formation. These results show that the MHM2 PrP carrying Lys, Ala, Trp, Arg or His at residue 218 binds to PPMF with a greater affinity than does wt MHM2 with Gln at 218. These findings also contend that the two polymorphic Hu residues Glu and Lys interact very differently with Mo PPMF. Mutant MHM2 Prp(E218) binds more weakly

to Mo PPMF than does wt MHM2 PrP(Q218) which results in MHM2 PrP(E218) not being converted into PrP^{Sc} and no inhibition of the conversion of wt MHM2 PrP^C into PrP^{Sc}. In contrast, mutant MHM2 PrP(K218) binds more tightly to Mo PPMF than does wt MHM2 PrP(Q218) which results in both MHM2 PrP(K218) not being converted into PrP^{Sc} and inhibition of the conversion of wt MHM2 PrP^C into PrP^{Sc}.

The substitution of Lys, Ala or Pro at residue 214 did not inhibit conversion of epitope tagged wt MHM2 PrP^C into PrP^{Sc}. In contrast, Ile, Glu or Trp at position 214 inhibited wt PrP^{Sc} formation. These results argue that the MHM2 PrP carrying Ile, Glu or Trp at position 214 binds to PPMF with a greater affinity than does wt MHM2 with Val at 218.

EXAMPLE 16

MUTATION AT Mo RESIDUE 216

MoPrP codon 216 which encodes a Gln residue in both Mo and Hu PrP was mutagenized because the side chain of this residue protrudes from the opposite face of the C-terminal α -helix at positions 214 and 218 and a mutation causes inherited prion disease. A Swedish family with GSS has been reported with a Q-R mutation, Hu codon 217 corresponds to Mo codon 216. Although brain sections showed PrP amyloid plaques, extracts showed neither infectivity nor protease resistant PrP^{Sc} on Western blots.

Arg was introduced at residue 216 in MHM2 PrP and MHM2 PrP(E218) which contains the Hu Glu residue at 218. Neither of these mutant PrPs acquired resistance when expressed in ScN2a cells. The Arg substitution at residue 216 inhibited conversion of epitope tagged wt MHM2 PrP^C into PrP^{Sc}. Whereas, no inhibition was observed if both residues 216 and 218 were mutated. These findings show that the Q-R mutation destabilizes the structure of PrP^C leading to inherited prion disease and prevents folding into a protease resistant PrP^{Sc} molecule. The Mo Glu residue at 218 allows MHM2 PrP(R216) to compete with wt MHM2 PrP for binding to PPMF whereas the Hu Glu residue decreases the affinity of this protein for PPMF. With the Hu Glu residue, no inhibition of the conversion of wt MHM2 PrP into PrP^{Sc} was observed.

EXAMPLE 17

MUTATIONS AT Mo RESIDUES 215, 221 AND 222

Substitution of Gln for Thr at MoPrP residue 215, Ala for Ser at Mo 221 or Lys for Glu at Mo 222 did not inhibit conversion of these epitope tagged MHM2 PrP molecules into PrP^{Sc} when expressed into PrP^{Sc} when expressed in ScN2a cells (Table 1). Co-expression of these mutant MHM2 PrPs with wt MHM2 PrP did not inhibit wt PrP^{Sc} formation. When the V214D mutation was introduced into MHM2 PrP(S221A), the protein was not converted into PrP^{Sc} suggesting that the V214D mutation prevented binding to PPMF.

10

EXAMPLE 18

MUTATION AT Mo RESIDUE 171

Results obtained show that mutations of Mo residues 214 and 218 inhibit PrP^{Sc} formation, showing that the surface of the C-terminal α -helix with the side chains of these residues is the binding site for PPMF. The NMR structure of SHa PrP90-231 shows a loop composed of residues 165-173 immediately adjacent to the PPMF binding site on the helix raising the possibility that one or more of these residues also participates in the binding to PPMF. To explore this possibility, we constructed mutant MHM2 PrP(N170S) and MHM2 PrP(Q171R) and transfected the DNAs into ScN2a cells. MHM2 PrP(N170S) was converted into PrP^{Sc} whereas MHM2 PrP(Q171R) was not. The Ser substitution at Mo residue 170 did not inhibit conversion of epitope tagged wt MHM2 PrP^C into PrP^{Sc}, whereas the Arg substitution inhibited PrP^{Sc} formation. These findings argue that Q171 in MoPrP forms a discontinuous epitope with V214 and Q218 to which PPMF binds.

EXAMPLE 19

SHA-PP INHIBITS CONVERSION MHM2 INTO PrP^{Sc}

The level of SHA-PP^C expression in Tg(SHA-PP)/Prp^{Sc} mice was directly proportional to the length of the incubation time after inoculation with Mo prions. To simulate these conditions in ScN2a cells, epitope-tagged MHM2 PrP was cocexpressed with SHA-PP or chimeric SHA/MoPrP. SHA-PP and chimeric MH3 PrP inhibited conversion of MHM2 PrP into PrP^{Sc} but this inhibition was relieved by substitution of Hu residues at positions 214, 218 and 219 designated HuA. As described above mutant PrP molecules with

Hu residues at 214 and 218 were not converted into PrP^{Sc} in ScN2a cells and did not inhibit the transformation of MHM2 PrP into PrP^{Sc}.

While neither SHa nor MH3 PrP^{Sc} could be efficiently converted into PrP^{Sc} in ScN2a cells reducing the extent of SHaPrP sequence resulted in a progressive increase in PrP^{Sc} formation. MHM2 PrP was converted into PrP^{Sc} at a level intermediate between MH3 PrP and MHM2 PrP. Expression of MHM2 PrP with MHM2 PrP decreased the level of conversion of MHM2 into PrP^{Sc}. Expression of MoPrP with MHM2 PrP did not inhibit the formation of PrP^{Sc} from MHM2 PrP.

The foregoing findings help define the order of addition during formation of the PPV PrP^{Sc}/PrP^{Sc} complex as well as the limits of central domain of PrP where PrP^{Sc} and PrP^{Sc} interact. When Hu residues at 214, 218 and 219 (HuA) were introduced into MHM2 or MH3 PrP, these chimeric PrPs no longer inhibited the conversion of MHM2 PrP into PrP^{Sc}. The HuA substitutions relieve inhibition by preventing the binding of the chimeric MH3HuA PrP. This shows that the lack of conversion of SHaPrP or MH3 PrP into PrP^{Sc} is not due to a low affinity for PPV. Instead, MoPrP^{Sc} does not stimulate conversion of SHaPrP or MH3 PrP into PrP^{Sc} even though these molecules are bound to PPV. This contention is supported by the ability of SHaPrP^{Sc} to stimulate conversion of SHaPrP into PrP^{Sc} in Tg(SHaPrP)PrP^{Sc} mice when MoPrP^{Sc} is coexpressed. These data show that PrP^{Sc} binds first to PPV and the protein PPV/PrP^{Sc} complex then binds to PrP^{Sc}.

The introduction of the HuA sequences into MH3 relieved the inhibition of MHM2 PrP conversion into PrP^{Sc}. This shows that the PPV binding site does not include SHa residues 203 and 205. Instead, these residues are part of the central domain where PrP^{Sc} and PrP^{Sc} interact because lack of conversion of MH3 into PrP^{Sc} by MoPrP^{Sc} in ScN2a cells could be partially overcome by changing the two SHaPrP and MH3 PrP are not converted into PrP^{Sc}, they are not released from PPV which in turn prevents MHM2 PrP from binding and being converted.

The methods and materials are the same as those described for Example 5 above.

EXAMPLE 20

ASSAY FOR COMPOUNDS INHIBITING PrP^{Sc} FORMATION

MATERIALS AND METHODS

Persistently infected scrapie mouse neuroblastoma cells (ScN2a) also expressing (MHM2) PrP^{Sc} were used in the assay. The ScN2a cells were grown and maintained at 37°C in MEM medium supplemented with 10% Fetal Bovine Serum or 1% lipid reduced serum for the positive controls using mevastatin which does inhibit PrP^{Sc} formation. Cells grown on 60 mm dishes were rinsed with PBS.

The monoclonal antibody 3F4 is raised against Syrian hamster (SHa) PrP 27-30. U.S. Patent 4,806,627, issued February 21, 1989, discloses monoclonal antibody 263K 3F4, produced by cell line ATCC HB9222 deposited on October 8, 1986, which is incorporated herein by reference. The cell line producing the antibody can be obtained from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852. To distinguish the signals of MHM2 constructs from endogenous mouse PrP^{Sc}, 3F4 mAb was used since 3F4 exclusively recognizes (SHa)PrP, as well as the Met 109 - Met 112 epitope of the MHM2 construct.

An anti-serum R073 is raised in a rabbit against SDS-PAGE purified (SHa) PrP 27-30 that reacts with SHa PrP, MoPrP and MHM2PrP. The R073 antibody is used to detect endogenous MoPrP in ScN2a cells.

Compounds screened in the assay of the invention are listed in the Compounds Table and were purchased from Sigma, Sigma-Aldrich For Rare Chemicals, NOVABiochem and CHMSRV-SQ. Solutions of the compounds are prepared and applied to the cell culture medium at a final concentration of 10 μ M.

SCREENING PROCEDURE

Final screening was carried out using ScN2a cells transfected with a pSPOX vector carrying the chimeric construct shown in Figures 1 and 2 using DOTAP Liposomal Transfection Reagent purchased from Boehringer-Mannheim. Twenty micrograms of plasmid is used for a 60 mm dish. Twenty four hours after transfection of the cells, compound is applied to the MEM medium at a final concentration of 10 μ M and incubated for 3 days. The cells were lysed four days after transfection. After the cells were lysed, they were (1)

subjected to Proteinase K digestion; (2) contacted with labeled antibodies; and (3) subjected to Western blot analysis.

Negative controls were run wherein the cells were incubated in the same manner but not exposed to any compound. Positive controls were run where the cells incubated in the same manner but were exposed to neovastatin.

Twenty nine compounds were tested. Twenty three did not inhibit PrP^{Sc} formation and three of the twenty three were cytotoxic. Six of the compounds did inhibit PrP^{Sc} formation and four of the six were cytotoxic. The results are shown in the Compound Table below with the best results obtained with Adenyl (3'-5') Cytidine and N-Phthaloyl-DL-Histidine in that both were not cytotoxic and did inhibit PrP^{Sc} formation.

COMPOUNDS TABLE

		SIGMA			
		Number	Name of Compound	Ref.	Solubility Screening Results
5	15 at room temperature		Bufoetin Disulfate Salt	B2845	Water 1 mg/ml Negative Not toxic
	16 at 4 degrees		Purimycin Antinomycinolide	P7130	Water 50 mg/ml Negative Not toxic
	17 at -20 degrees		2'-3'-Di-O-Acetylguanoside	D2847	Dimethyl Formamide Negative Not toxic
	18 at room temperature		Purine Riboside	P9278	Water 50 mg/ml Positive Cytotoxic
	19 at 4 degrees		Gonadotropin	G1279	Water 50 mg/ml Negative Not toxic
	20 at -20 degrees		pClu-Oly-Arg-Phe-Amide	P9799	Dimethyl Formamide Negative Not toxic
	21 at 4 degrees		Ribostamycin	R2255	Water 50 mg/ml Negative Not toxic
	22 at 4 degrees		Dibekacin	D7024	Water 50 mg/ml Positive Cytotoxic
	23 at 4 degrees		Sisomycin	S8757	Water 50 mg/ml Test again
	25 at room temperature		P-Adenylguanosyl (2'-5'-Adenosine-3'-Inosine) Riboside	A8675	Water 10 mg/ml (Cytidine) Negative Not toxic
15	26 at -20 degrees		6-(1-Hydroxyguanylinosine) Riboside	H5272	Water 50 mg/ml Test again
	27 at -20 degrees		Adenyl(3'-5')Cytidine	A7435	Water 10 mg/ml Positive Not toxic
	28 at -20 degrees		Guanyl(2'-5')Guanosine	G1133	Water 10 mg/ml Negative Not toxic
	22 at -20 degrees		2'-3'-Di-O-Acetylguanosine	D6626	Dimethyl Formamide Positive Cytotoxic
	33 at -20 degrees		Guanosine 2'- & 3'-Monophosphate	G8002	Water Negative Not toxic
	34 at 4 degrees		MTH-DL-Ariginine HCl	M3756	Water 50 mg/ml Positive Cytotoxic
	35 at -20 degrees		Adenosine 5'-Carboxylic Acid	A0895	NH ₄ OH 30 mg/ml Negative Not toxic
	36 at -20 degrees		Inosine 3'-Monophosphate	I7628	Water 50 mg/ml Negative Not toxic
	38 at 4 degrees		Allantoinic Acid	A7753	NH ₄ OH 30 mg/ml Negative Cytotoxic
	39 at 4 degrees		Amikacin	A3650	Water 50 mg/ml Negative Not toxic
20	40 at -20 degrees		2'-O-Acetylguanosyladenosine 3'-5'-Cyclic Monophosphate	A6688	Water 50 mg/ml Positive Cytotoxic

CLAIMS

That which is claimed is:

1. An isolated composition of Prion Protein Modulator Factor (PPMF) defined by the following characteristics:
 - (a) binding to PrP^{Sc} and facilitating a conformational change from PrP^C to PrP^{Sc};
2. The PPMF of claim 1, further characterized by:
 - (b) does not bind to PrP^{Sc};
 - (c) is species specific thereby binding only to PrP^C of a genetically similar species;
 - (d) acting as rate limiting component in conversion of PrP^C to PrP^{Sc};
 - (e) binding of PPMF not effected by a residue at position 216, 217, 220, 221 or 222 in human PrP^C and a corresponding position in a different species;
 - (f) facilitating formation of PrP^C/PrP^{Sc} complexes;
 - (g) being released from PrP^C upon its conversion to PrP^{Sc};
 - (h) diminishing the activation energy barrier between PrP^C and PrP^{Sc} and thereby facilitating formation of PrP^{Sc};
 - (i) having its binding effected by residues at position 215 and 219 in hamster PrP^C and corresponding positions in a different species; and
 - (j) preferably further characterized by:
 - (i) having its binding to PrP^C effected by a residue at position 171 in sheep PrP^C and a corresponding position in a different species;
 - (k) binding to a discontinuous epitope on PrP^C which epitope comprises residues 172, 215 and 219 in human PrP^C and corresponding residues in different species; and
 - (l) being encoded by a nucleotide sequence of a species selected from the group consisting of human, pig, cow, sheep, mouse and hamster.
 3. A mimetic of a phosphore of an epitope on PrP^C which binds PPMF which mimetic preferably has a surface which matches the geometric specifications of any of figures 4, 12, 13, 14 or 15.

5

SIGMA-ALDRICH FOR RARE CHEMICALS				
Number	Name of Compound	Ref.	Solubility	Screening Results
44 at room temperature	1,1-Thiois 3,5-Dimethyl hydantoin (C10H11ANOAS1) M.W. 286 g/mol	S94, 973-6	Water	Negative Non toxic
45 at room temperature	N Phthaloyl-DL Histidine (C14H11NO4) M.W. 285 g/mol	S40, 985-5	NaOH IN	Positive Non toxic
46 at room temperature	7-acetoxymethyl 5 benzamido hexah 7 methyl 5 oximidazo (C21H25NO6) M.W. 415 g/mol	S76, 001-3	DMF	Negative Non toxic
47 at room temperature	3 nitro 4,4' methylendianiline (C13H13NO2) M.W. 253 g/mol	S59, 207-6	Methanol	Negative Non toxic
48 at room temperature	n-ethylbenzoyl-L-Histidinol (C13H15NO2) M.W. w45 g/mol	S93, 331-7	Methanol	Negative Non toxic

NOVA Biochem				
Number	Name of Compound	Ref.	Solubility	Screening Results
49 at -20 degrees	BOC-Val-Gly-Ala-Phe-Asp-Asp M.W. 555.7 g/mol	03-324512	DMF (50%) and water	Negative Non toxic

20 While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

25

4. A method of inhibiting Prp^{as} formation, comprising:

administering a mimetic pharmacophore of PPMF which binds an epitope on Prp^c wherein the pharmacophore preferably has a surface which matches a negative image of any of figures 4, 12, 13, 14 or 15 and more preferably is selected from the group consisting of Adenyl(3'5')Cytidine, Purine Riboside, Dibekacin, N Phthaloyl-DL-Histidine, 2'3'-Di-O-Acetyladenosine, and MTH-DL-Arginine.

5. A Prp gene characterized by having a native wild type codon replaced with a non-native codon at a position which encodes an amino acid making up a portion of a binding epitope to PPMF.

6. The Prp gene of claim 5, wherein the gene encodes a Prp protein selected from the group consisting of:

a protein which binds PPMF more tightly than native Prp binds PPMF to the extent that Prp^{as} formation is blocked;

a Prp protein which does not bind PPMF sufficiently will to allow for the formation of Prp^{as}.

wherein the gene is selected from the group consisting of:

a cow Prp gene and the wild type codon replaced is from a position selected from the group consisting of 171, 175, 218 and 222;

a sheep Prp gene and the wild type codon replaced is from a position selected from the group consisting of 171, 175, 218 and 222;

a mouse Prp gene and the wild-type codon replaced is from a position selected from the group consisting of 167, 171, 214 and 218;

a human Prp gene and the wild-type codon replaced is from a position selected from the group consisting of 167, 171, 214 and 218; and

a gene wherein the replaced codons are in the Prp gene at positions which encode amino acids in the range of 160-180 and 210-225.

7. An isolated nucleotide sequence encoding the PPMF of claim 1.

8. An antibody selected from the group consisting of:

an antibody immunospecific for the PPMF of claim 1 which antibody binds PPMF at its PPMF/Prp^c binding site;

a Prp^c antibody that binds Prp^c at its epitope for binding the PPMF of claim 1; and

a Prp^c antibody that binds Prp^c at an epitope which prevents the formation of a

5 Prp^c/PPMF/Prp^{as} complex.

9. An animal resistant to prion disease which animal is selected from the group consisting of a mouse, a hamster, a human, a pig, a sheep and a cow wherein Prp genes of the animal have wild-type codons substituted with different codons at two or more codon positions which encode an amino acid of an epitope of Prp^c which binds PPMF;

preferably wherein four native codon positions which encode an amino acid of an epitope of Prp^c which binds PPMF are substituted with non-native codons;

further wherein the animal is preferably selected from the group consisting of:

a cow and the wild-type codons replaced are from a position selected from the group consisting of 171, 175, 218 and 222;

a sheep and the wild-type codons replaced are from a position selected from the group consisting of 171, 175, 218 and 222;

a mouse and the wild-type codons replaced are from a position selected from the group consisting of 167, 171, 214 and 218; and

a mouse and the wild-type codons replaced are from a position selected from the group consisting of 167, 171, 214 and 218.

10. A method of identifying a compound which inhibits Prp^{as} formation, comprising:

25 providing a culture of cells which cells are genetically engineered to express Prp^c protein;

adding Prp^{as} to the cells;

adding a compound to the cells;

lysing the cells after allowing for sufficient time for converting Prp^c to Prp^{as}; and

analyzing to determine an effect on Prp^{as} formation.




FIG. 1




FIG. 2




FIG. 11



3/17

Mo	MH2	MH3	MH4	MH5	MH6	MH7	MH8	MH9	MH10	MH11	MH12	MH13	MH14	MH15	MH16	MH17	MH18	MH19	MH20	MH21	MH22	MH23	MH24	MH25	MH26	MH27	MH28	MH29	MH30	MH31	MH32	MH33	MH34	MH35	MH36	MH37	MH38	MH39	MH40	MH41	MH42	MH43	MH44	MH45	MH46	MH47	MH48	MH49	MH50	MH51	MH52	MH53	MH54	MH55	MH56	MH57	MH58	MH59	MH60	MH61	MH62	MH63	MH64	MH65	MH66	MH67	MH68	MH69	MH70	MH71	MH72	MH73	MH74	MH75	MH76	MH77	MH78	MH79	MH80	MH81	MH82	MH83	MH84	MH85	MH86	MH87	MH88	MH89	MH90	MH91	MH92	MH93	MH94	MH95	MH96	MH97	MH98	MH99	MH100	MH101	MH102	MH103	MH104	MH105	MH106	MH107	MH108	MH109	MH110	MH111	MH112	MH113	MH114	MH115	MH116	MH117	MH118	MH119	MH120	MH121	MH122	MH123	MH124	MH125	MH126	MH127	MH128	MH129	MH130	MH131	MH132	MH133	MH134	MH135	MH136	MH137	MH138	MH139	MH140	MH141	MH142	MH143	MH144	MH145	MH146	MH147	MH148	MH149	MH150	MH151	MH152	MH153	MH154	MH155	MH156	MH157	MH158	MH159	MH160	MH161	MH162	MH163	MH164	MH165	MH166	MH167	MH168	MH169	MH170	MH171	MH172	MH173	MH174	MH175	MH176	MH177	MH178	MH179	MH180	MH181	MH182	MH183	MH184	MH185	MH186	MH187	MH188	MH189	MH190	MH191	MH192	MH193	MH194	MH195	MH196	MH197	MH198	MH199	MH200	MH201	MH202	MH203	MH204	MH205	MH206	MH207	MH208	MH209	MH210	MH211	MH212	MH213	MH214	MH215	MH216	MH217	MH218	MH219	MH220	MH221	MH222	MH223	MH224	MH225	MH226	MH227	MH228	MH229	MH230	MH231	MH232	MH233	MH234	MH235	MH236	MH237	MH238	MH239	MH240	MH241	MH242	MH243	MH244	MH245	MH246	MH247	MH248	MH249	MH250	MH251	MH252	MH253	MH254	MH255	MH256	MH257	MH258	MH259	MH260	MH261	MH262	MH263	MH264	MH265	MH266	MH267	MH268	MH269	MH270	MH271	MH272	MH273	MH274	MH275	MH276	MH277	MH278	MH279	MH280	MH281	MH282	MH283	MH284	MH285	MH286	MH287	MH288	MH289	MH290	MH291	MH292	MH293	MH294	MH295	MH296	MH297	MH298	MH299	MH300	MH301	MH302	MH303	MH304	MH305	MH306	MH307	MH308	MH309	MH310	MH311	MH312	MH313	MH314	MH315	MH316	MH317	MH318	MH319	MH320	MH321	MH322	MH323	MH324	MH325	MH326	MH327	MH328	MH329	MH330	MH331	MH332	MH333	MH334	MH335	MH336	MH337	MH338	MH339	MH340	MH341	MH342	MH343	MH344	MH345	MH346	MH347	MH348	MH349	MH350	MH351	MH352	MH353	MH354	MH355	MH356	MH357	MH358	MH359	MH360	MH361	MH362	MH363	MH364	MH365	MH366	MH367	MH368	MH369	MH370	MH371	MH372	MH373	MH374	MH375	MH376	MH377	MH378	MH379	MH380	MH381	MH382	MH383	MH384	MH385	MH386	MH387	MH388	MH389	MH390	MH391	MH392	MH393	MH394	MH395	MH396	MH397	MH398	MH399	MH400	MH401	MH402	MH403	MH404	MH405	MH406	MH407	MH408	MH409	MH410	MH411	MH412	MH413	MH414	MH415	MH416	MH417	MH418	MH419	MH420	MH421	MH422	MH423	MH424	MH425	MH426	MH427	MH428	MH429	MH430	MH431	MH432	MH433	MH434	MH435	MH436	MH437	MH438	MH439	MH440	MH441	MH442	MH443	MH444	MH445	MH446	MH447	MH448	MH449	MH450	MH451	MH452	MH453	MH454	MH455	MH456	MH457	MH458	MH459	MH460	MH461	MH462	MH463	MH464	MH465	MH466	MH467	MH468	MH469	MH470	MH471	MH472	MH473	MH474	MH475	MH476	MH477	MH478	MH479	MH480	MH481	MH482	MH483	MH484	MH485	MH486	MH487	MH488	MH489	MH490	MH491	MH492	MH493	MH494	MH495	MH496	MH497	MH498	MH499	MH500	MH501	MH502	MH503	MH504	MH505	MH506	MH507	MH508	MH509	MH510	MH511	MH512	MH513	MH514	MH515	MH516	MH517	MH518	MH519	MH520	MH521	MH522	MH523	MH524	MH525	MH526	MH527	MH528	MH529	MH530	MH531	MH532	MH533	MH534	MH535	MH536	MH537	MH538	MH539	MH540	MH541	MH542	MH543	MH544	MH545	MH546	MH547	MH548	MH549	MH550	MH551	MH552	MH553	MH554	MH555	MH556	MH557	MH558	MH559	MH560	MH561	MH562	MH563	MH564	MH565	MH566	MH567	MH568	MH569	MH570	MH571	MH572	MH573	MH574	MH575	MH576	MH577	MH578	MH579	MH580	MH581	MH582	MH583	MH584	MH585	MH586	MH587	MH588	MH589	MH590	MH591	MH592	MH593	MH594	MH595	MH596	MH597	MH598	MH599	MH600	MH601	MH602	MH603	MH604	MH605	MH606	MH607	MH608	MH609	MH610	MH611	MH612	MH613	MH614	MH615	MH616	MH617	MH618	MH619	MH620	MH621	MH622	MH623	MH624	MH625	MH626	MH627	MH628	MH629	MH630	MH631	MH632	MH633	MH634	MH635	MH636	MH637	MH638	MH639	MH640	MH641	MH642	MH643	MH644	MH645	MH646	MH647	MH648	MH649	MH650	MH651	MH652	MH653	MH654	MH655	MH656	MH657	MH658	MH659	MH660	MH661	MH662	MH663	MH664	MH665	MH666	MH667	MH668	MH669	MH670	MH671	MH672	MH673	MH674	MH675	MH676	MH677	MH678	MH679	MH680	MH681	MH682	MH683	MH684	MH685	MH686	MH687	MH688	MH689	MH690	MH691	MH692	MH693	MH694	MH695	MH696	MH697	MH698	MH699	MH700	MH701	MH702	MH703	MH704	MH705	MH706	MH707	MH708	MH709	MH710	MH711	MH712	MH713	MH714	MH715	MH716	MH717	MH718	MH719	MH720	MH721	MH722	MH723	MH724	MH725	MH726	MH727	MH728	MH729	MH730	MH731	MH732	MH733	MH734	MH735	MH736	MH737	MH738	MH739	MH740	MH741	MH742	MH743	MH744	MH745	MH746	MH747	MH748	MH749	MH750	MH751	MH752	MH753	MH754	MH755	MH756	MH757	MH758	MH759	MH760	MH761	MH762	MH763	MH764	MH765	MH766	MH767	MH768	MH769	MH770	MH771	MH772	MH773	MH774	MH775	MH776	MH777	MH778	MH779	MH780	MH781	MH782	MH783	MH784	MH785	MH786	MH787	MH788	MH789	MH790	MH791	MH792	MH793	MH794	MH795	MH796	MH797	MH798	MH799	MH800	MH801	MH802	MH803	MH804	MH805	MH806	MH807	MH808	MH809	MH810	MH811	MH812	MH813	MH814	MH815	MH816	MH817	MH818	MH819	MH820	MH821	MH822	MH823	MH824	MH825	MH826	MH827	MH828	MH829	MH830	MH831	MH832	MH833	MH834	MH835	MH836	MH837	MH838	MH839	MH840	MH841	MH842	MH843	MH844	MH845	MH846	MH847	MH848	MH849	MH850	MH851	MH852	MH853	MH854	MH855	MH856	MH857	MH858	MH859	MH860	MH861	MH862	MH863	MH864	MH865	MH866	MH867	MH868	MH869	MH870	MH871	MH872	MH873	MH874	MH875	MH876	MH877	MH878	MH879	MH880	MH881	MH882	MH883	MH884	MH885	MH886	MH887	MH888	MH889	MH890	MH891	MH892	MH893	MH894	MH895	MH896	MH897	MH898	MH899	MH900	MH901	MH902	MH903	MH904	MH905	MH906	MH907	MH908	MH909	MH910	MH911	MH912	MH913	MH914	MH915	MH916	MH917	MH918	MH919	MH920	MH921	MH922	MH923	MH924	MH925	MH926	MH927	MH928	MH929	MH930	MH931	MH932	MH933	MH934	MH935	MH936	MH937	MH938	MH939	MH940	MH941	MH942	MH943	MH944	MH945	MH946	MH947	MH948	MH949	MH950	MH951	MH952	MH953	MH954	MH955	MH956	MH957	MH958	MH959	MH960	MH961	MH962	MH963	MH964	MH965	MH966	MH967	MH968	MH969	MH970	MH971	MH972	MH973	MH974	MH975	MH976	MH977	MH978	MH979	MH980	MH981	MH982	MH983	MH984	MH985	MH986	MH987	MH988	MH989	MH990	MH991	MH992	MH993	MH994	MH995	MH996	MH997	MH998	MH999	MH1000	MH1001	MH1002	MH1003	MH1004	MH1005	MH1006	MH1007	MH1008	MH1009	MH1010	MH1011	MH1012	MH1013	MH1014	MH1015	MH1016	MH1017	MH1018	MH1019	MH1020	MH1021	MH1022	MH1023	MH1024	MH1025	MH1026	MH1027	MH1028	MH1029	MH1030	MH1031	MH1032	MH1033	MH1034	MH1035	MH1036	MH1037	MH1038	MH1039	MH1040	MH1041	MH1042	MH1043	MH1044	MH1045	MH1046	MH1047	MH1048	MH1049	MH1050	MH1051	MH1052	MH1053	MH1054	MH1055	MH1056	MH1057	MH1058	MH1059	MH1060	MH1061	MH1062	MH1063	MH1064	MH1065	MH1066	MH1067	MH1068	MH1069	MH1070	MH1071	MH1072	MH1073	MH1074	MH1075	MH1076	MH1077	MH1078	MH1079	MH1080	MH1081	MH1082	MH1083	MH1084	MH1085	MH1086	MH1087	MH1088	MH1089	MH1090	MH1091	MH1092	MH1093	MH1094	MH1095	MH1096	MH1097	MH1098	MH1099	MH1100	MH1101	MH1102	MH1103	MH1104	MH1105	MH1106	MH1107	MH1108	MH1109	MH1110	MH1111	MH1112	MH1113	MH1114	MH1115	MH1116	MH1117	MH1118	MH1119	MH1120	MH1121	MH1122	MH1123	MH1124	MH1125	MH1126	MH1127	MH1128	MH1129	MH1130	MH1131	MH1132	MH1133	MH1134	MH1135	MH1136	MH1137	MH1138	MH1139	MH1140	MH1141	MH1142	MH1143	MH1144	MH1145	MH1146	MH1147	MH1148	MH1149	MH1150	MH1151	MH1152	MH1153	MH1154	MH1155	MH1156	MH1157	MH1158	MH1159	MH1160	MH1161	MH1162	MH1163	MH1164	MH1165	MH1166	MH1167	MH1168	MH1169	MH1170	MH1171	MH1172	MH1173	MH1174	MH1175	MH1176	MH1177	MH1178	MH1179	MH1180	MH1181	MH1182	MH1183	MH1184	MH1185	MH1186	MH1187	MH1188	MH1189	MH1190	MH1191	MH1192	MH1193	MH1194	MH1195	MH1196	MH1197	MH1198	MH1199	MH1200	MH1201	MH1202	MH1203	MH1204	MH1205	MH1206	MH1207	MH1208	MH1209	MH1210	MH1211	MH1212	MH1213	MH1214	MH1215	MH1216	MH1217	MH1218	MH1219	MH1220	MH1221	MH1222	MH1223	MH1224	MH1225	MH1226	MH1227	MH1228	MH1229	MH1230	MH1231	MH1232	MH1233	MH1234	MH1235	MH1236	MH1237	MH1238	MH1239	MH1240	MH1241	MH1242	MH1243	MH1244	MH1245	MH1246	MH1247	MH1248	MH1249	MH1250	MH1251	MH1252	MH1253	MH1254	MH1255	MH1256	MH1257	MH1258	MH1259	MH1260	MH1261	MH1262	MH1263	MH1264	MH1265	MH1266	MH1267	MH1268	MH1269	MH1270	MH1271	MH1272	MH1273	MH1274	MH1275	MH1276	MH1277	MH1278	MH1279	MH1280	MH1281	MH1282	MH1283	MH1284	MH1285	MH1286	MH1287	MH1288	MH1289	MH1290	MH1291	MH1292	MH1293	MH1294	MH1295	MH1296	MH1297	MH1298	MH1299	MH1300	MH1301	MH1302	MH1303	MH1304	MH1305	MH1306	MH1307	MH1308	MH1309	MH1310	MH1311	MH1312	MH1313	MH1314	MH1315	MH1316	MH1317	MH1318	MH1319	MH1320	MH1321	MH1322	MH1323	MH1324	MH1325	MH1326	MH1327	MH1328	MH1329	MH1330	MH1331	MH1332
----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------

5/17

WO 98/55132


PCT/US98/10104



SUBSTITUTE SHEET (RULE 26)

SUBSTITUTE SHEET (RULE 26)

8/17



SUBSTITUTE SHEET (RULE 26)

SUBSTITUTE SHEET (RULE 26)




FIG. 7

9/17

10/17




FIG. 8




FIG. 9

12/17




FIG. 12




FIG. 10

13/17




FIG. 13

14/17




FIG. 14

15/17




FIG. 15

16/17





FIG. 16

17/17



INTERNATIONAL SEARCH REPORT		International application No. PCT/US98/0104
A. CLASSIFICATION OF SUBJECT MATTER IPC(O) : A61K 3406; C07K 100; C07H 2102, 2104 US CL : 530/240, 336, 336/23.1 According to International Patent Classification (IPC) or to both national classification and IPC		
B. PIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 330/240, 336, 336/23.1		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base used, where practicable, search terms used) Please See Extra Sheet		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	KANEKO, K. et al., Evidence for protein X binding to a discontinuous epitope on the cellular prion protein during scrapie prion propagation. Proceedings of the National Academy of Science, USA, September 1997, Vol. 94, pages 10069-10074, see entire document.	1, 2, and 7
Y	TELLING, G.C. et al., Prion Propagation in Mice Expressing Human and Chimera PrP Transgenes Implicates the Interaction of Cellular PrP with Another Protein. Cell. 06 October 1995, Vol. 83, pages 79-90, see entire document.	1, 2, and 7
Y,P	PRUSINER, S.B. et al., Prion Protein Biology. Cell. 01 May 1998, Vol. 93, pages 337-348, see entire document.	1, 2, and 7
<input checked="" type="checkbox"/> Further documents are listed in the continuation of this U. <input type="checkbox"/> See parent family agent.		
* Special categories of cited documents: "X" prior document published after the international filing date of priority date and not in conflict with the application but cited to inform the applicant of the state of the art "Y" document published on or after the international filing date and in conflict with the application but cited to inform the applicant of the state of the art "Y,P" document published on or after the international filing date and in conflict with the application but cited to inform the applicant of the state of the art "L" document published on or after the international filing date and in conflict with the application but cited to inform the applicant of the state of the art "O" document published on or after the international filing date and in conflict with the application but cited to inform the applicant of the state of the art "P" document published prior to the international filing date but later than the priority date of the application		
Date of the final completion of the international search		Date of mailing of the international search report
18 AUGUST 1998		07 OCT 1998
Name and mailing address of the ISAUS International Search Authority Box PCT Washington, D.C. 20231 Fax: 202/335-3230		Authorized officer BRIAN K. HANNA Telephone No. (703) 304-0196

INTERNATIONAL SEARCH REPORT		International application No.
C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		PCT/US98/0104
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	TELLING, G.C. et al., Transmission of Creutzfeldt-Jakob disease from humans to transgenic mice expressing chimeric human-mouse prion protein. Proceedings of the National Academy of Sciences, USA, October 1994, Vol. 91, pages 9936-9940, see entire document.	1, 2, and 7
Y	COHEN, F.E. et al., Structural Clues to Prion Replication, Science, 22 April 1994, Vol. 264, pages 530 and 531, see entire document.	1, 2, and 7.
Y.P	YEHIELY, F. et al., Identification of candidate proteins binding to prion proteins. Neurobiology of Disease, 1997, Vol. 3, No. 4, pages 330-355, see entire document.	1, 2, and 7

Form PCT/ISA210 (continuation of first sheet(!)) July 1997) 8

INTERNATIONAL SEARCH REPORT	
<div> <div>International Application No.</div> <div>PC703910104</div> </div>	
<div>Box I</div> <div>Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)</div>	<div>This international report has not been established in report of certain claims under Article 17(2)(a) for the following reasons:</div>
<div>1. <input type="checkbox"/> Claims Nos.:</div> <div>because they relate to subject matter not required to be searched by this Authority, namely:</div>	
<div>2. <input type="checkbox"/> Claims Nos.:</div> <div>because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:</div>	
<div>3. <input type="checkbox"/> Claims Nos.:</div> <div>because they are dependent claims and are not included in accordance with the second and third sentences of Rule 6.4(a).</div>	
<div>Box II</div> <div>Observations where validity of inventions is lacking (Continuation of Item 2 of first sheet)</div>	<div>This International Searching Authority found multiple inventions in this international application, as follows:</div> <div>Please See Exam Sheet.</div>
<div>1. <input type="checkbox"/> As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.</div>	
<div>2. <input type="checkbox"/> As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.</div>	
<div>3. <input type="checkbox"/> As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:</div>	
<div>4. <input checked="" type="checkbox"/> No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:</div> <div>1, 2, and 7</div>	
<div>Remark on Patent</div> <div><input type="checkbox"/> <input type="checkbox"/></div>	<div>The additional search fees were accompanied by the applicant's protest.</div> <div>No protest accompanied the payment of additional search fees.</div>

Form PCT/ISA210 (continuation of first sheet(!)) July 1997) 8

PCT/US99/10104

Electronic data bases consulted (Name of data base and where practicable terms used):

Search Terms: price?; protein?; modular?; pruniser?/au; ocobes?/au; protein x

This ISA found multiple investments as follows:

Group I, claim(s) 1, 2, and 7, drawn to prior art modulator factors (PPMF)

THIS PAGE BLANK (USPTO)